arXiv:0904.1519v1 [math.AG] 9 Apr 2009

ELLIPTIC K3 SURFACES WITH ABELIAN AND DIHEDRAL GROUPS OF SYMPLECTIC AUTOMORPHISMS

ALICE GARBAGNATI

Abstract. We analyze K3 surfaces admitting an elliptic fibration $\mathcal{E}$ and a finite group G of symplectic automorphisms preserving this elliptic fibration. We construct the quotient elliptic fibration $\mathcal{E}/G$ comparing its properties to the ones of $\mathcal{E}$.
We show that if $\mathcal{E}$ admits an n-torsion section, its quotient by the group of automorphisms induced by this section admits again an n-torsion section.
Considering automorphisms coming from the base of the fibration, we can describe the Mordell–Weil lattice of a fibration described by Kloosterman. We give the isometries between lattices described by the author and Sarti and lattices described by Shioda and by Griees and Lam. Moreover we show that for certain groups H of G, H subgroups of G, a K3 surface which admits H as group of symplectic automorphisms actually admits G as group of symplectic automorphisms.

0. Introduction

An automorphism on a K3 surface is called symplectic if it leaves invariant the nowhere vanishing holomorphic two form of the K3 surface. Nikulin was the first to study this kind of automorphisms. One of the main properties of a finite group G of symplectic automorphisms on a K3 surface X is that the desingularization of the quotient X/G is again a K3 surface. It is a general fact (cf. [N1], [W]) that if X is a K3 surface admitting a finite group G as group of symplectic automorphisms, the lattice $\Omega_G := (H^2(X,\mathbb{Z})^G)^\perp$ is primitively embedded in $NS(X)$. Similarly, if Y is a K3 surface obtained as desingularization of X/G, there exists a lattice, called M_G, (which contains the curves arising from the desingularization of X/G) which is primitively embedded in $NS(Y)$.

Here we analyze examples of elliptic K3 surfaces admitting symplectic automorphisms preserving the elliptic fibration, and we describe the quotient surfaces.
In [GS1] and [GS2] elliptic fibrations on K3 surfaces are considered to analyze some properties of the family of K3 surfaces admitting finite abelian groups of symplectic automorphisms. In particular, if a K3 surface X admits an elliptic fibration $\mathcal{E}$ with a torsion section t, it admits a symplectic automorphism σ_t, which is the translation by the torsion section, and which acts as the identity on the base of the fibration. This automorphism preserves the fibration, hence, on the desingularization of the quotient surface X/σ_t, there is an elliptic fibration, induced by the one on X. In *Section 4*, Proposition 4.1, we give our main result about such elliptic fibrations: if $\mathcal{E}$ is the generic elliptic fibration admitting a certain group $G=\langle t\rangle$ (generated by a torsion section t) as Mordell–Weil group, then the elliptic fibrations $\mathcal{E}/\sigma_t$ and $\mathcal{E}$ have isomorphic Mordell–Weil groups and the same type of singular fibers. This implies that the Néron–Severi groups of the surface X and $\widetilde{X/\sigma_t}$ are isometric and we determine them. Using the fact that $\widetilde{X/\sigma_t}$ is obtained as quotient of a K3 surface by a group of symplectic automorphisms, we show that its Néron–Severi group is isometric to $U\oplus M_G$. In [GS1] and [GS2] the Néron–Severi group of X (and hence the one of $\widetilde{X/\sigma_t}$ since they are isometric) was described considering its relation with the lattice Ω_G, and in fact $NS(X)$ was obtained as overlattice of finite index of $U(n)\oplus\Omega_G$, where $n=|G|$.
The relation between the elliptic fibration with a certain torsion section and the elliptic fibration obtained as quotient of this by the automorphism induced by section can be analyzed in a more general

---

2000 Mathematics Subject Classification: 14J28, 14J50, 14J10.
Key words: K3 surfaces, elliptic fibrations, symplectic automorphisms, lattices.

setting. In *Section 2* we recall that if E is an elliptic curve defined over a field $\mathbb{K}$ with a rational n-torsion point T_n, and there is a primitive n-th root of unity in $\mathbb{K}$, then the elliptic curve $E/\langle T_n\rangle$ has again a rational n-torsion point. In the particular case of the elliptic fibration with base $\mathbb{P}^1$ (analyzed in *Section* 3) this implies that the quotient of the elliptic fibration with the automorphism induced by an n-torsion section, is again an elliptic fibration with an n-torsion section.

Until now, we considered automorphisms of elliptic fibrations with base $\mathbb{P}^1$ (and in particular on K3 surfaces) which act trivially on the base. But, of course, it is possible to construct also automorphisms of the elliptic fibrations which are induced by automorphisms of the base of the fibrations. This kind of automorphisms is interesting because they are related to base changes of elliptic fibrations. In *Section 5* we construct elliptic fibrations with base $\mathbb{P}^1$ admitting certain automorphisms acting also on the basis and we compute the equation of the elliptic surface obtained as quotient of the elliptic fibrations by these automorphisms. In *Section 6* we come back to the case of the K3 surfaces and we consider the automorphisms constructed in Section 5, which are symplectic in the case of K3 surface. The main result of this part is that we can construct elliptic fibrations such that the Néron–Severi group is $U\oplus\Omega_G$ for certain abelian groups G (this is in certain sense the analogue of result of Proposition 4.1, where we constructed surfaces with Néron–Severi $U\oplus M_G$). Moreover, this construction gives a very nice interpretation of the lattice Ω_G, which appears in this situation as the Mordell–Weil lattice of the fibration. As a consequence, we proved that certain lattices found by Shioda in [Shio3] and [Shio4] are isometric to the lattice $\Omega_{\mathbb{Z}/n\mathbb{Z}}$ for $n=5,6$ computed in [GS1] and [GS2].
Moreover, considering automorphisms which come from the base of the fibration, we can construct some dihedral groups of symplectic automorphisms on K3 surface. Let $\mathcal{D}_n$ be the dihedral group of order $2n$. By [X], if $\mathcal{D}_n$ appears as group of symplectic automorphisms of a K3 surface, then $n=3,4,5,6$. In *Section 7* a family of elliptic K3 surfaces admitting the group $\mathcal{D}_4$ as group of symplectic automorphisms is described. The automorphisms generating $\mathcal{D}_4$ came from automorphisms of the base of the fibration. The family considered here was described in [Kl] to give an example of an elliptic K3 surface with rank of Mordell–Weil group equal to 15. However in [Kl] the Mordell–Weil lattice and the Néron–Severi group of this fibration are not described. Here we prove that the Mordell–Weil lattice has to be equal to $\Omega_{\mathcal{D}_4}$ and, considering a special member of the family, we compute explicitly this lattice and we prove that it is isometric to a lattice described in [GL]. Moreover, (as in case of the abelian group G) we prove that the Néron–Severi group has to be isometric to $U\oplus\Omega_{\mathcal{D}_4}$.

In *Section 8* we consider examples of elliptic K3 surfaces admitting the dihedral groups $\mathcal{D}_n$ as groups of symplectic automorphisms. We consider both automorphisms which come from automorphisms of the base, and which are induced by torsion section, so in a certain sense we combine the constructions considered before. The examples considered in this section are used to prove an interesting phenomenon, which was already described in [G2] for the group of symplectic automorphisms $\mathcal{D}_5$: let H be a subgroup of G. If G appears as group of symplectic automorphisms on a K3 surface, of course so does H. For certain groups H and G, if one constructs the family of the K3 surfaces admitting H as group of symplectic automorphisms, one finds that all the members of this family admit the group G as group of symplectic automorphisms. We prove that this happens for the pairs (H,G) $(\mathbb{Z}/n\mathbb{Z},\mathcal{D}_n)$, $n=5,6$, $(\mathbb{Z}/2\mathbb{Z}\times\mathbb{Z}/4\mathbb{Z},\mathbb{Z}/2\mathbb{Z}\times\mathcal{D}_4)$, $(\mathbb{Z}/3\mathbb{Z}\times\mathbb{Z}/3\mathbb{Z},\mathfrak{A}_{3,3})$. In particular this implies that for each of these pairs $\Omega_H\simeq\Omega_G$. In *Section* 9 we reconsider the problem analyzed in Section 8 in a more general setting. We consider pairs of groups (H,G) such that H is a subgroup of G and such that the K3 surfaces admitting H as group of symplectic automorphisms admits in fact the group G as groups of symplectic automorphisms and we show that this property is equivalent to a property related only to the lattices Ω_H and Ω_G (Proposition 9.1). We prove that there are pairs (H,G) with this property which are not among the ones described in Section 8, but the ones described in Section 8 are the only ones such that $\mathrm{rk}\Omega_H$=16.

I would like to thank Bert van Geemen for his invaluable help, Jaap Top for useful discussions and Tetsuji Shioda for the lectures he held in Milan, which aided me greatly in working on this paper.

1. Background material

Here we collect the main results on symplectic automorphisms on K3 surfaces and on elliptic fibrations on $\mathbb{P}^1$ which will be useful in the following.

1.1. **Symplectic automorphisms on K3 surfaces.**

Definition 1.1. *An automorphism σ on a K3 surface X is symplectic if σ^* acts as the identity on $H^{2,0}(X)$, that is $\sigma^*(\omega_X)=\omega_X$, where ω_X is a nowhere vanishing holomorphic two form on X. Equivalently σ is symplectic if the isometry induced by σ^* on the transcendental lattice is the identity.*

Let σ be an automorphism of finite order on a K3 surface. The desingularization of X/σ is a K3 surface if and only if σ is symplectic.
The main results about the finite abelian groups of symplectic automorphisms on K3 surfaces were obtained by Nikulin in [N1]. In particular he classifies these groups and proves that the isometries induced on the second cohomology group of the K3 surfaces, called Λ_{K3}, by a finite abelian group of symplectic automorphisms on a K3 surface are essentially unique ([N1, Definition 4.6, Theorem 4.7]). As a consequence of this theorem one obtains that if a K3 surface X admits a certain group G as group of symplectic automorphisms, then the lattice $\Omega_G:=(\Lambda_{K3}^G)^\perp$, which is primitively embedded in $NS(X)\subset H^2(X,\mathbb{Z})$ (cf. [N1], [GS1], [GS2]), does not depend on X, but only on G. The lattices Ω_G are computed in [GS1], [GS2] for each possible finite abelian group G of symplectic automorphisms on a K3 surface.
Hence, the main result about the finite abelian groups of symplectic automorphisms is:

Proposition 1.1. ([N1], [GS1], [GS2]) *Let G be a finite abelian group. A K3 surface X admits G as group of symplectic automorphisms if and only if Ω_G is primitively embedded in $NS(X)$.*

In [N1] the number and the kind of the singularities of the quotient of a K3 surface by a finite abelian group of symplectic automorphisms are classified. If X is a K3 surface admitting a finite abelian group of symplectic automorphisms G, then X/G has singularities of type A_l. Hence in the desingularization $\widetilde{X/G}$ we obtain some rational curves M_i which are the exceptional curves of the blow ups which desingularize the surface.

Definition 1.2. *Let M_G be the minimal primitive sublattice of $NS(\widetilde{X/G})$ containing the curves M_j arising from the resolution of the singularities of X/G.*

The lattice M_G contains some linear combinations of the M_j with rational, non integer, coefficients. For each finite abelian group G which can act symplectically on a K3 surface a description of the lattice M_G is given in [N1, Section 7].

In [Mu] and in [X] the finite (but not necessary abelian) group of symplectic automorphisms are classified. In [X] the singularities of the surface X/G (where G is a finite group of symplectic automorphisms of X) are described. Moreover the main result of [N1] (i.e. the uniqueness of the isometry induced on the second cohomology group by the finite abelian group of symplectic automorphisms) holds also in the non abelian case, [W]. Hence Proposition 1.1 holds also for finite groups G, not necessary abelian. The lattice Ω_G has not yet been explicitly determined for each G.

1.2. **Elliptic fibration on $\mathbb{P}^1$.**

Definition 1.3. *Let X be a surface and B a curve. An elliptic fibration $\mathcal{E}:X\rightarrow B$ on the surface X is a fibration such that the generic fiber is a smooth curve of genus one and such that a section $s:B\rightarrow X$ is fixed. We call this section the zero section.*

We recall that every elliptic fibration can be regarded as an elliptic curve over the function field of the basis. We will assume $B\simeq\mathbb{P}^1$. Under this assumption each elliptic fibration admits a minimal Weierstrass equation of the form

$$y^2=x^3+A(\tau,\sigma)x+B(\tau,\sigma),\quad A(\tau,\sigma),\ B(\tau,\sigma)\in\mathbb{C}[\tau,\sigma]_{hom},\quad \deg A(\tau,\sigma)=4m,\ \deg B(\tau,\sigma)=6m$$

for a certain $m \in \mathbb{N}_{>0}$ and where there exists no a polynomial $C(\tau,\sigma)$ such that $C(\tau,\sigma)^4|A(\tau,\sigma)$ and $C(\tau,\sigma)^6|B(\tau,\sigma)$.
We use the notation $A(\tau)$ and $B(\tau)$ to indicate the polynomial $A(\tau,1)$ and $B(\tau,1)$.
If $m=1$, then X is a rational surface, if $m=2$, X is a K3 surface, if $m>2$ X the Kodaira dimension of X, $k(X)$, is 1. More in general $h^{2,0}(X)=dimH^{2,0}(X)=m-1$ (and a basis of $H^{2,0}(X)$ is given by $\tau^i dx/y\wedge d\tau$, $i=0,\ldots,m-2$).
Hence in particular if $\deg A(\tau)\leq 4$ and $\deg B(\tau)\leq 6$, then the surface X is rational.

There are finitely many singular fibers. The possible singular fibers on an elliptic fibration are described by Kodaira (cf. for example [Mi]). In particular the fibers of type I_n, $n>2$ are made up of n rational curves meeting as a polygon with n edges. We will call C_0 the irreducible component of a reducible fiber which meets the zero section. The irreducible components of a fiber of type I_n are called C_i where $C_i\cdot C_{i+1}=1$ and $i\in\mathbb{Z}/n\mathbb{Z}$. Under the assumption $C_0\cdot s=1$, these conditions identify the components completely once the component C_1 is chosen, so these conditions identify the components up to the transformation $C_i\leftrightarrow C_{-i}$ for all $i\in\mathbb{Z}/n\mathbb{Z}$. The components of a reducible fiber of type I_n have multiplicity one, so a section can intersect a fiber of type I_n in any component.

The set of the sections of an elliptic fibration form a group (the Mordell–Weil group), with the group law which is induced by the one on the fibers. Let r be the rank of the Mordell–Weil group (recall that if there are no sections of infinite order then $r=0$) and $\rho=\rho(X)$ be the Picard number of the surface X and Red be the set $Red=\{v\in\mathbb{P}^1\mid F_v \text{ is reducible}\}$. Then

$$\rho(X)=rkNS(X)=r+2+\sum_{v\in Red}(m_v-1)$$

(cfr. [Shio1, Section 7]) where m_v is the number of irreducible components of the fiber F_v.

Definition 1.4. *The* **trivial lattice** Tr_X *(or* Tr*) of an elliptic fibration on a surface* X *is the lattice generated by the class of the fiber, the class of the zero section and the classes of the irreducible components of the reducible fibers which do not intersect the zero section.*

The lattice Tr admits the hyperbolic plane U as sublattice and its rank is $rk(Tr)=2+\sum_{v\in Red}(m_v-1)$.

Theorem 1.2. [Shio1, Theorem 1.3] *The Mordell–Weil group of the elliptic fibration on the surface* X *is isomorphic to the quotient* $NS(X)/Tr=:E(K)$.

In Section 8 of [Shio1] a pairing on $E(K)$ is defined. The value of this pairing on a section P depends only on the intersection between the section P and the reducible fibers and between P and the zero section. Now we recall the definition and the properties of this pairing.
Let $E(K)_{tor}$ be the set of the torsion elements in the group $E(K)$.

Lemma 1.3. [Shio1, Lemma 8.1, Lemma 8.2] *For any* $P\in E(K)$ *there exists a unique element* $\phi(P)$ *in* $NS(X)\otimes\mathbb{Q}$ *such that:*
i) $\phi(P)\equiv(P)$ *mod* $Tr\otimes\mathbb{Q}$ *(where* (P) *is the class of* P *modulo* $Tr\otimes\mathbb{Q}$*)*
ii) $\phi(P)\perp Tr$.
The map $\phi: E(K)\rightarrow NS(X)\otimes\mathbb{Q}$ *defined above is a group homomorphism such that* $Ker(\phi)=E(K)_{tor}$.

Theorem 1.4. [Shio1, Theorem 8.4] *For any* $P,Q\in E(K)$ *let* $\langle P,Q\rangle=-\phi(P)\cdot\phi(Q)$ *(where* $\cdot$ *is induced on* $NS(X)\otimes\mathbb{Q}$ *by the cup product). This defines a symmetric bilinear pairing on* $E(K)$*, which induces the structure of a positive definite lattice on* $E(K)/E(K)_{tor}$.
In particular if $P\in E(K)$*, then* P *is a torsion section if and only if* $\langle P,P\rangle=0$.
For any $P,Q\in E(K)$ *the pairing* $\langle -,-\rangle$ *is*

$$\begin{array}{rcl}\langle P,Q\rangle & = & \chi+P\cdot s+Q\cdot s-P\cdot Q-\sum_{v\in Red}\mathrm{contr}_v(P,Q)\\ \langle P,P\rangle & = & 2\chi+2(P\cdot s)-\sum_{v\in Red}\mathrm{contr}_v(P)\end{array}$$

where χ *is the holomorphic Euler characteristic of the surface and the rational numbers* $\mathrm{contr}_v(P)$ *and* $\mathrm{contr}_v(P,Q)$ *are given in the table below*

$$\begin{array}{llllll} & I_2 & I_n & I_n^* & IV^* & III^* \\ \mathrm{contr}_v(P) & 2/3 & i(n-i)/n & \begin{cases} 1 \text{ if } i=1 \\ 1+n/4 \text{ if } i=n-1 \text{ or } i=n \end{cases} & 4/3 & 3/2 \\ \mathrm{contr}_v(P,Q) & 1/3 & i(n-j)/n & \begin{cases} 1/2 \text{ if } i=1 \\ 2+n/4 \text{ if } i=n-1 \text{ or } i=n \end{cases} & 2/3 & - \end{array} \tag{1}$$

where the numbering of the fibers is the one described before, P *and* Q *meet the fiber in the component* C_i *and* C_j *and* $i \leq j$.

The pairing defined in the theorem is called the **height pairing**. This pairing will be used to determine the intersection of the torsion sections of the elliptic fibrations with the irreducible components of the reducible fibers.
The Mordell–Weil group equipped with the height pairing is called Mordell–Weil lattice. We remark that if $rk(Tr) = 2$ (i.e. there are no reducible fibres and hence no torsion sections), the height pairing on the Mordell–Weil group is the restriction of the cup product on $NS(X)$ to $\phi(E(K))$. So, if $rk(Tr) = 2$, the Mordell–Weil lattice is the sublattice $\phi(E)$ of $NS(X)$.

2. Quotient of elliptic curves by groups generated by torsion points

An elliptic curve is a pair (E, O), where E is a cubic curve defined over a field $\mathbb{K}$ and O is a rational point on this curve. We will assume $char\mathbb{K} \neq 2, 3$. Each elliptic curve admits an equation of the following form

$$y^2 = f_3(x), \quad \text{where } f_3(x) \in \mathbb{K}[x], \quad deg(f_3) = 3 \text{ and } \Delta(f_3) \neq 0,$$

where Δ is the discriminant of a polynomial f. The point $O = (0 : 1 : 0)$ with respect to the projective coordinates $(x : y : z)$ is a point on the elliptic curve.
We recall the following result:

Proposition 2.1. *Assume that* $\zeta_n \in \mathbb{K}$ *is a primitive* n*-th root of unity. Let* E_n *be an elliptic curve, defined over the field* $\mathbb{K}$*, admitting an* n*-torsion rational point* Q*.*
Then the elliptic curve $E_n/\langle Q\rangle$ *has an* n*-torsion rational point* R*. Moreover the elliptic curve* $(E_n/\langle Q\rangle)/\langle R\rangle$ *is isomorphic to the curve* E_n*.*

Proof. Let us consider the group $E_n[n] = \{S \in E_n(\overline{\mathbb{K}}) \text{ such that } nS = 0\}$.
For each $\sigma \in Gal(\overline{\mathbb{K}}/\mathbb{K})$ and $S \in E_n[n]$, also $\sigma(S) \in E_n[n]$. So $Gal(\overline{\mathbb{K}}/\mathbb{K})$ acts on $E_n[n] \simeq \mathbb{Z}/n\mathbb{Z} \times \mathbb{Z}/n\mathbb{Z}$. Thus we get an homomorphism $\rho_n : Gal(\overline{\mathbb{K}}/\mathbb{K}) \to Aut(E_n[n]) \simeq GL_2(\mathbb{Z}/n\mathbb{Z})$.
Let Q and $\widetilde{R}$ be two generators of $E[n]$. By hypothesis Q is a rational point and so for each $\sigma \in Gal(\overline{\mathbb{K}}/\mathbb{K})$, $\sigma(Q) = Q$. Hence $\rho_n(\sigma) = \begin{bmatrix} 1 & \alpha_\sigma \\ 0 & \beta_\sigma \end{bmatrix}$.
Let $e_n : E_n[n] \times E_n[n] \to \mu_n = \{n\text{-th roots of unity}\}$ be the Weil pairing. It induces an isomorphism $\wedge^2 E_n[n] \to \mu_n$. Since σ acts on $E_n[n]$, $\wedge^2(\sigma)$ acts on $\wedge^2 E_n[n]$. The map $\wedge^2(\sigma)$ is the multiplication by $det(\rho_n(\sigma))$. As $\zeta_n \in \mathbb{K}$, $det(\rho_n(\sigma)) = 1$. This implies that $\beta_\sigma = 1$ for each $\sigma \in Gal(\overline{\mathbb{K}}/\mathbb{K})$. So

$$\rho_n(\sigma) = \begin{bmatrix} 1 & \alpha_\sigma \\ 0 & 1 \end{bmatrix} \text{ and } \sigma(\widetilde{R}) = \alpha_\sigma Q + \widetilde{R} \text{ for each } \sigma \in Gal(\overline{\mathbb{K}}/\mathbb{K}).$$

The morphism

$$\phi : E_n \to E_n/\langle Q\rangle$$

is defined over $\mathbb{K}$ and thus $R := \phi(\widetilde{R})$ is a rational n-torsion point on $E/\langle Q\rangle$. The composition $E_n \to E_n/\langle Q\rangle \to (E_n/\langle Q\rangle)/\langle R\rangle$ is a multiplication by n on E_n. □

For certain values of n the curve E_n, which admits an n-torsion rational point, the quotient map ϕ and of the curve $E_n/\langle Q\rangle$ can be given explicitly. For $n = 2$ these equations are very well known (see. e. g. [ST, Pag. 79]), for $n = 3$ they can be found in [T]. As example here we give a similar result for $n = 4$.

Proposition 2.2. *The curve E admits a 4-torsion rational point if and only if, up to a change of coordinate, it admits an equation of the form*

$$y^2 = x(x^2 + (e^2 - 2f)x + f^2).$$

Let us assume that E admits such an equation, then $Q = (f, ef)$ is a 4-torsion rational point. Let $G = \langle Q \rangle$.
The curve E/G has equation

$$y^2 = x(x^2 - 2(e^2 + 4f)x + (e^2 - 4f)^2).$$

If $i \in \mathbb{K}$ with $i^2 = -1$, then the equation becomes $y^2 = x(x^2 + (h^2 - 2g)x + g^2)$ with $g = 4f - e^2$, $h = 2ie$. In particular if $i \in \mathbb{K}$, E/G admits a 4-torsion rational point, which is the image of a 4-torsion point on $E(\bar{\mathbb{K}})$ not in G.
The quotient map is:

$$\phi(P) = \begin{cases} \left(\frac{(f+x)^2y^2}{(f-x)^2x^2}, \frac{y(f+x)((f-x)^4-4e^2fx^2)}{x^2(f-x)^3}\right) & \textit{if } P = (x,y) \neq (0,0), (f, ef), (f, -ef), O_E \\ O_{E/G} & \textit{if } P = (x,y) = (0,0), (f, ef), (f, -ef), O_E \end{cases}$$

Proof. The elliptic curve E has in particular a point of order two, so has an equation of the form $y^2 = x(x^2 + ax + b)$ (see e.g. [ST, pag. 79]). To determine a and b we observe that a smooth elliptic curve of the fibration has a point Q of order four if $Q + Q = R = (0, 0)$ which is the rational point of order 2. Geometrically this means that the tangent to the elliptic curve through Q must intersect the curve exactly in the point R. Taking $Q = (e, ef)$ the equation is

$$y^2 = x(x^2 + (e^2 - 2f)x + f^2).$$

To obtain the equation of the quotient curve E/G, we will consider the quotient by the point of order two R, and next the quotient by the image of Q on the curve $E/\langle R \rangle$. The point R corresponds to $(0, 0)$. By [ST, pag. 79] the curve $E/\langle R \rangle$ is

$$y^2 = x(x^2 - 2(e^2 - 2f)x + ((e^2 - 2f)^2 - 4f^2))$$

and so

$$y^2 = x(x - e^2)(x - e^2 + 4f).$$

This curve has three 2-torsion rational points: $(0, 0)$, $(e^2, 0)$, $(e^2 - 4f, 0)$. The 4-torsion point Q is mapped to the point $\bar{Q} = (e^2, 0)$ on $E/\langle R \rangle$, which is clearly a 2-torsion point on $E/\langle R \rangle$. We consider a transformation which sends $\bar{Q}$ to $(0, 0)$. Under this transformation the equation becomes

$$y^2 = (x + e^2)x(x + 4f).$$

Its quotient by the point $(0, 0)$ (i.e. by the point $\bar{Q}$) is

$$y^2 = x(x^2 - 2(e^2 + 4f)x + (e^2 - 4f)^2)$$

If $i \in \mathbb{K}$, then the equation becomes

$$y^2 = x(x^2 + (h^2 - 2g)x + g^2) \quad \text{with} \quad g = 4f - e^2,\ h = 2ie$$

which is the equation of an elliptic curve with a rational 4-torsion point (g, gh).
To describe explicitly the map we have to consider the map given in [ST, Pag. 79], to compose them with the translation $(x, y) \mapsto (x - e^2, y)$ on $E/\langle R \rangle$, and to compose again with the map given in [ST, Pag. 79]. □

3. Automorphisms induced by sections

Now we consider an elliptic fibration $\mathcal{E} : X \to \mathbb{P}^1$. We will indicate with $\mathcal{E}$ also the equation of the fibration viewed as elliptic curve over $k(\mathbb{P}^1)$. If the equation is in the standard Weierstrass form, we can assume that there is a zero section which is the map $\tau \mapsto (0 : 1 : 0)$. The other sections of the elliptic fibration $\mathcal{E}$, if any, induce automorphisms on X.

Definition 3.1. *If t is a section, then we call σ_t the automorphism defined as translation by t. It acts as the identity on the basis of the fibration $\mathbb{P}^1$.*
The group $G = \langle\sigma_t\rangle$ is a group of automorphisms of X which preserves the elliptic fibration $\mathcal{E}$. Let us consider the desingularization, $\widetilde{X/G}$, of the quotient X/G. Since the elliptic fibration $\mathcal{E}$ is preserved by G, $\widetilde{X/G}$ admits an elliptic fibration induced by $\mathcal{E}$. We call it $\mathcal{E}/G$.

The torsion sections of the fibration $\mathcal{E} : X \to \mathbb{P}^1$ are torsion rational points of $\mathcal{E}$ considered as an elliptic curve over the field $k(\mathbb{P}^1)$.
If the section t is an n-torsion section, the automorphisms σ_t is of order n. In particular, if $G = torsMW(\mathcal{E})$, then G is a finite abelian group of automorphisms on X.

Remark 3.1. Let t be a section of order n on X. If X is a rational surface, then $2 \leq n \leq 6$; if X is a K3 surface, then $2 \leq n \leq 8$. Indeed, if t is a torsion section, then the height formula computed on t is zero (Theorem 1.4). This implies that there are a certain number and type of reducible fibers. The Picard number of a rational (resp. K3) elliptic surface is 10 (resp. at most 20), this gives a bound for the order of the possible torsion sections on these surfaces. □

The following lemmas describe explicitly the action of the automorphism σ_t on the semistable reducible fibers of the fibration. The case of instable fibers is similar, but is not useful in the following.

Lemma 3.2. [GS2, Lemma 2.2] *Let t be an n-torsion section and F_v be a reducible fiber of type I_d of the fibration and let us suppose that the section t meets the fiber F_v in the component C_i.*
Let r be the minimal number such that $(\sigma_t)^r_{|F_v}(C_j) = C_j$, $\forall j \in \mathbb{Z}/d\mathbb{Z}$.
If $\gcd(d,n) = 1$ then $r = 1$, $i = 0$ and $\sigma_t(C_j) = C_j$, $\forall j \in \mathbb{Z}/d\mathbb{Z}$.
If $\gcd(d,n) \neq 1$ and $i = 0$, then $r = 1$ and $\sigma_t(C_j) = C_j$, $\forall j \in \mathbb{Z}/d\mathbb{Z}$.
If $\gcd(d,n) \neq 1$ and $i \neq 0$, then $r = d/\gcd(d,i)$, $r|\gcd(d,n)$, and $\sigma_t(C_j) = C_{j+i}$.

Lemma 3.3. *Let $\mathcal{E}$ be a fibration with a singular fiber F_v of type I_d and an n-torsion section t as in the previous lemma and let $G = \langle\sigma_t\rangle$. Then the image of the fiber F_v on $\mathcal{E}/G$ is a fiber of type $I_{\frac{dn}{r^2}}$.*

Proof. The singular fiber F_v/G has $\frac{d}{r}$ component meeting as a polygon (because $(\sigma_t)_{|F_v}$ identifies each component with r other components). Since σ_t^r fixes the points of intersection of the components in F_v (and hence the stabilizer of each point is $\mathbb{Z}/\frac{n}{r}\mathbb{Z}$), each vertex of the polygon F_v/G is a singular point of type $A_{\frac{n}{r}-1}$. So in $\widetilde{X/G}$ there is a tree of $\frac{n}{r}-1$ rational curves over such a point. Hence the fiber corresponding to F_v on $\mathcal{E}/G$ is a polygon with $\frac{d}{r}\cdot\frac{n}{r}$ edges, i.e. a fiber of type $I_{\frac{dn}{r^2}}$. □

Proposition 3.4. *Let $\mathcal{E}_G$ be an elliptic fibration on $\mathbb{P}^1$ admitting either $G = \mathbb{Z}/m\mathbb{Z}$ or $G = \mathbb{Z}/m\mathbb{Z} \times \mathbb{Z}/n\mathbb{Z}$ as torsion part of the Mordell–Weil group. Then $G \subset tors(MW(\mathcal{E}_G/G))$.*

Proof. This follows by Proposition 2.1 and by the fact that $\zeta_n \in k(\mathbb{P}^1)$ for each n. □

The equations of elliptic fibrations $\mathcal{E}_G$ for $G = \mathbb{Z}/n\mathbb{Z}$, $n = 2,3,4,5,6,8$ are the ones given in [GS2, Table 1] with different degrees for the polynomials with variable τ. If S is the surface admitting the fibration $\mathcal{E}_G$ and $h^{2,0}(S) = m-1$, then the degrees of the polynomial are obtained multiply the degrees given in [GS2] by $m/2$. If the number obtained is not an integer, there exists no a surface S such that $h^{2,0}(S) = m-1$ and S admits an elliptic fibration of with an elliptic fibration with that group as torsion part of the Mordell–Weil group. In particular the elliptic fibrations $\mathcal{E}_G$ with $G \subset MW(\mathcal{E}_G)$ are parameterized by the coefficients of the polynomials, and hence by an irreducible variety.

Proposition 3.5. *Let $\mathcal{E}_G$ be a generic elliptic fibration on $\mathbb{P}^1$ admitting $G = \mathbb{Z}/n\mathbb{Z}$, $n = 2,3,4,5,6,8$ as torsion part of the Mordell–Weil group (here generic means that the coefficients of the polynomials in τ have been chosen generically). Then the singular fibers of $\mathcal{E}_G$ and of $\mathcal{E}_G/G$ are of the same type. Moreover $\mathcal{E}_G$ and $\mathcal{E}_G/G$ have isomorphic Néron–Severi group.*

Proof. The type of the singular fibers of an elliptic fibration can be completely determined by the zeros of the polynomials $A(\tau)$, $B(\tau)$ and $\Delta(\tau)$ and hence can be directly computed by the equation

on $\mathcal{E}_G$.
We will consider as example the case $n = 6$, the others are very similar. The singular fibers of an elliptic fibration with a 6-torsion section t are $mI_6 + mI_3 + mI_2 + mI_1$ where $\deg k(\tau) = \deg l(\tau) = m$ with the notation of [GS2]. We deduce the intersection between the singular fibers and the section t by the height formula (cf. Theorem 1.4). Indeed the only possibility to have a 6-torsion section (and hence a section with height pairing equal to zero) is that t meets all the reducible fibers in the component C_1. By Lemma 3.3, this is enough to describe the elliptic fibration $\mathcal{E}_G/G$ on $\widetilde{X/\sigma_t}$. Indeed the fibers of type I_6 (resp. I_3, I_2, I_1) on X correspond to fibers of type I_1 (resp. I_2, I_3, I_6). Hence the elliptic fibration $\mathcal{E}_G/G$ has as singular fibers $mI_6 + mI_3 + mI_2 + mI_1$.
The Néron–Severi group of an elliptic fibration is generated by the class of the fiber, the classes of the sections and the classes of the reducible components of the reducible fibers. The hypothesis on the generality of the elliptic fibration $\mathcal{E}_G$ guarantees that $MW(\mathcal{E}_G) = MW(\mathcal{E}_G/G) = G$. So $\mathcal{E}_G$ and $\mathcal{E}_G/G$ have the "same" sections, and the "same" singular fibers and hence they have the "same" Néron–Severi group. □

4. Elliptic K3 surfaces and automorphisms induced by sections

In this section we focalize our attention on K3 surfaces.
The automorphisms σ_t induced on an elliptic K3 surface X by n-torsions section are symplectic. Indeed on X we have the nowhere vanishing holomorphic 2-form $dx/y \wedge d\tau$, where dx/y is a 1-form on the elliptic curve in the fiber. Since σ_t acts trivially on the base of the fibration and as a translation on the fibers (hence fixing the 1-forms on it), $dx/y \wedge d\tau$ is fixed by σ_t.

Proposition 4.1. *Let X_G be a K3 surface admitting the elliptic fibration $\mathcal{E}$ with $tors(MW(\mathcal{E})) = G$. Then $G \subset tors(MW(\mathcal{E}/G))$.*
Moreover, if X_G is such that $\mathcal{E}$ is the generic elliptic fibration with $tors(MW(\mathcal{E})) = G$, then $NS(X_G) \simeq NS(\widetilde{X_G/G}) \simeq U \oplus M_G$.

Proof. The fact that $G \subset tors(MW(\mathcal{E}))$ is Proposition 3.4.
The possible G such that there exists an elliptic fibration $\mathcal{E}$ on a K3 surface with $torsMW(\mathcal{E}) = G$ are $G = \mathbb{Z}/n\mathbb{Z}$, $2 \leq n \leq 8$, $G = (\mathbb{Z}/n\mathbb{Z})^2$ with $n = 2, 3, 4$, $G = \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/n\mathbb{Z}$ with $n = 4, 6$. For the group $\mathbb{Z}/n\mathbb{Z}$ with $n = 2, 3, 4, 5, 6, 8$ the fact that in the generic case $NS(X_G) \simeq NS(\widetilde{X_G/G})$ is consequence of Proposition 3.5. For the groups $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/m\mathbb{Z}$ with $m = 4, 6$ the fact that in the generic case $NS(X_G) \simeq NS(\widetilde{X_G/G})$ can be proved as in Proposition 3.5, considering the equation given in [GS2]. The groups $(\mathbb{Z}/n\mathbb{Z})^2$ are the full n-torsion group of the elliptic curve $\mathcal{E}$ (which is defined over $k(\mathbb{P}^1)$). The quotient $E/E[n]$ corresponds to a multiplication by n on the curve, and hence gives an isomorphic curve. This of course implies that $NS(X_G) \simeq NS(\widetilde{X_G/G})$.
It remains the case $G = \mathbb{Z}/7\mathbb{Z}$. There exists only one possible elliptic fibration admitting a 7-torsion section. By [Shim] this elliptic fibration has to admit 3 fibers of type I_7 and 3 fibers of type I_1 and it has no sections of infinte order (because its trivial lattice has rank 20 which is the maximal possible Picard number for a K3 surface). By Lemma 3.3, $\mathcal{E}/G$ admits 3 fibers of type I_7 (corresponding to the fibers of type I_1 on $\mathcal{E}$) and 3 fibers of type I_1 (corresponding to the fibers of type I_7 on $\mathcal{E}$). Moreover, by Proposition 3.4, such an elliptic fibration admits a 7-torsion section. Hence, as in Proposition 3.5, $NS(X_G) \simeq NS(\widetilde{X_G/G})$.
Let us prove that $NS(\widetilde{X_G/G}) \simeq U \oplus M_G$ in the case $G = \mathbb{Z}/5\mathbb{Z}$. The proof in the other cases is essentially the same. In this case the trivial lattice is $U \oplus A_4^{\oplus 4}$. The singular fibers of this fibration are $4I_5 + 4I_1$. The elliptic fibration $\mathcal{E}/G$ has $4I_1 + 4I_5$ as singular fibers (cf. Lemma 3.3). In $\mathcal{E}/G$ the components ($C_i^{(j)}$, $i, j = 1, \ldots, 4$) of the fibers I_5 which do not meet the zero section are the rational curves arising by the resolution of the four singularities of type A_4 on X_G/G (cf. proof of Lemma

3.3). They are the curves M_i of Definition 1.2 and hence by [N1] the class

$$v=\frac{1}{5}\left[\sum_{i=1}^{2}(4C_1^{(i)}+3C_2^{(i)}+2C_3^{(i)}+C_4^{(i)})+\sum_{j=3}^{4}(3C_1^{(j)}+6C_2^{(j)}+4C_3^{(j)}+2C_4^{(j)})\right]$$

is a class in $NS(\widetilde{X_G/G})$. The irreducible components which do not intersect the zero section of the reducible fibers of elliptic fibration $\mathcal{E}/G$ and the class v generate the lattice M_G. They are orthogonal to the class of the fiber (because they are linear combinations of components of a fiber) and to the class of the zero section, hence $NS(\widetilde{X_G/G})\hookleftarrow U\oplus M_G$ with finite index. By definition M_G is primitive in $NS(\widetilde{X_G/G})$ and since U is a unimodular lattice there are no overlattice of $U\oplus M_G$ such that M_G is primitive in them. So $NS(\widetilde{X_G/G})\simeq U\oplus M_G$. □

Remark 4.2. The class v in the proof of the previous proposition generates a copy of $\mathbb{Z}/5\mathbb{Z}$ in $NS(\widetilde{X_G/G})/Tr_{\widetilde{X_G/G}}\simeq MW(\widetilde{X_G/G})$ and this shows again that in $\mathcal{E}/G$ there is a 5-torsion section (without using Proposition 3.4). The class $t_1=s+2F-v$ is a class in $NS(\widetilde{X_G/G})$ corresponding to a 5-torsion section of the fibration. □

In case $G=\mathbb{Z}/2\mathbb{Z}$ Proposition 4.1 was already proved in [vGS, Proposition 4.2].

The K3 surfaces X_G are examples of K3 surfaces which both admit a finite abelian group G of symlectic automorphisms and are, at the same time, the desingularization of the quotient of a K3 surface by the same group of automorphisms. We observe that this is not the case for a generic K3 surface with a symplectic group of automorphisms, i.e. in general a K3 which admits a group G as finite abelian group of symplectic automorphisms cannot be obtained as quotient of another K3 surface by the same group of automorphisms.

Remark 4.3. Since $\widetilde{X_G/G}$ admits an elliptic fibration with G as torsion part of the Mordell–Weil group, it admits G as group of symplectic automorphisms, hence we can consider the surface Z_G, the desingularization of the quotient of $\widetilde{X_G/G}$ by G. The surface Z_G is isomorphic to X_G by the last statement in Proposition 2.1. □

5. AUTOMORPHISMS INDUCED BY AUTOMORPHISMS OF THE BASIS.

Until now we considered automorphisms which leave the base of the elliptic fibration invariant. Now we consider automorphisms which act also on the base of the fibration. Let us consider an automorphism π of $\mathbb{P}^1$. It induces the map $\nu:(x,y,\tau)\mapsto(x,y,\pi(\tau))$. In general this is not an automorphism of the fibration, (in general it does not preserve the fibration), but under some conditions (which depend on π) on the polynomials $A(\tau)$ and $B(\tau)$, it is.
A family of examples can be constructed considering the automorphisms of $\mathbb{P}^1$ $\pi_n:\tau\mapsto\zeta_n\tau$, where ζ_n is a primitive n-root of the unit. So $\nu_n:(x,y,\tau)\mapsto(x,y,\zeta_n\tau)$.

Let X be a surface with elliptic fibration $\mathcal{E}:y^2=x^3+A(\tau)x+B(\tau)$.
If $A(\pi_n(\tau))=A(\tau)$ and $B(\pi_n(\tau))=B(\tau)$, then ν_n is an automorphism of the X. The conditions on $A(\tau)$ and $B(\tau)$ imply that, $A(\tau)$ and $B(\tau)$ are actually polynomials in τ^n, so $A(\tau)=\sum_{i=0}^{l}a_i\tau^{in}$ and $B(\tau)=\sum_{i=0}^{h}b_i\tau^{in}$ and $\deg A(\tau)=nl$, $\deg B(\tau)=nh$. The automorphism ν_n fixes the two fibers on the point $\tau=0$ and $\tau=\infty$ (i.e. over the fixed points for π on $\mathbb{P}^1$). The quotient surface, $Y=X/\nu$, is smooth, and admits an elliptic fibration with equation $y^2=x^3+(\sum_{i=0}^{l}a_it^i)x+\sum_{i=0}^{h}b_it^i$.
Conversely given Y, X is obtained from Y by the base change $\tau\mapsto t=\tau^n$ (cf. [Mi, VI.4]). If the polynomials $A(\tau)=\sum_{i=0}^{l}a_i\tau^{in}$ and $B(\tau)=\sum_{i=0}^{h}b_i\tau^{in}$ are such that there does not exist a polynomial $C(\tau)$ such that $C(\tau)^4|A(\tau)$, $C(\tau)^6|B(\tau)$, then the equation obtained by the base change

is the Weierstrass form of the elliptic fibration $\mathcal{E}$ on the surface X. In fact we have the following commutative diagram

$$\begin{tikzcd}
X \arrow[r] \arrow[d] & Y = X/\nu_n \arrow[d] \\
\mathbb{P}^1_\tau \arrow[r] & \mathbb{P}^1_t = \mathbb{P}^1_\tau/\nu
\end{tikzcd}$$

We observe that the degree of the polynomials in t of the elliptic fibration on X/ν_n are l and h and the degree of the corresponding polynomials in τ on X are nl and nh. If X for example is an elliptic K3 surface and $n = 2$ (i.e. $4 < 2l < 9$ and $6 < 2h < 13$), then $Y = X/\nu_n$ is a rational elliptic surface (because $l \leq 4$ and $h \leq 6$). Hence it is possible that the Kodaira dimensions of X and X/ν_n are not the same. In particular if X is a K3 surface, X/ν_n is not. In fact the automorphisms ν_n do not fix the two holomorphic form $dx/y \wedge d\tau$ on X and hence they are not symplectic.

We now construct maps σ_n which act on $\mathbb{P}^1$ as π_n, but which act also on the fibers in such a way that one obtains symplectic automorphisms if the surface X is a K3 surface.

Definition 5.1. *Let σ_n be the map $\sigma_n : (x, y, \tau) \mapsto (\zeta_n^2 x, \zeta_n^3 y, \zeta_n \tau)$;*
μ_2 be the map $\mu_2 : (x, y, \tau) \mapsto (x, y, \frac{1}{\tau})$ and
ς_2 be the map $\varsigma_2 : (x, y, \tau) \mapsto (\frac{x}{\tau^4}, -\frac{y}{\tau^6}, \frac{1}{\tau})$.

Proposition 5.1. *Let $\mathcal{E}_n$ be an elliptic fibration with minimal equation $y^2 = x^3 + A(\tau)x + B(\tau)$. The map σ_n is an automorphism of the elliptic fibration $\mathcal{E}_n$ if and only if*

$$A(\tau) = \zeta_n^{-4} A(\zeta_n \tau) \text{ and } B(\tau) = \zeta_n^{-6} B(\zeta_n \tau). \tag{2}$$

If (2) holds, then the elliptic fibration $\mathcal{E}_n$ has equation:

$$y^2 = x^3 + \left(\sum_{i=-[\frac{4}{n}]}^{h} a_i \tau^{4+ni} \right) x + \sum_{i=-[\frac{6}{n}]}^{k} b_i \tau^{6+ni} \tag{3}$$

and the elliptic fibration $\mathcal{E}_n/\sigma_n$ has equation:

$$Y^2 = X^3 + \left(\sum_{i=-[\frac{4}{n}]}^{h} a_i T^{4+i} \right) X + \sum_{i=-[\frac{6}{n}]}^{k} b_i T^{6+i}.$$

In particular by the minimality of the equation of $\mathcal{E}_n$ it follows that $n \leq 6$.

Proof. Applying the map σ_n to $\mathcal{E}_n$, one obtains $y^2\zeta_n^6 = x^3\zeta_n^6 + A(\zeta_n\tau)x\zeta_n^2 + B(\zeta_n\tau)$. It is easy to check that it is again the equation of $\mathcal{E}_n$ if and only if (2) holds. The equation of the quotient surface is computed by multiplying the equation of the elliptic fibration $\mathcal{E}_n$ by τ^{6n-6} and considering $X := \tau^{2n-2}x$, $Y := \tau^{3n-3}y$, $T := \tau^n$ as coordinates of the quotient. The condition on n comes from the fact that if $n \geq 6$, then $\tau^4 | A(\tau)$ and $\tau^6 | B(\tau)$. □

Remark 5.2. The elliptic fibration $\mathcal{E}_n$ is obtained from the elliptic fibration $\mathcal{E}_n/\sigma_n$ by a base change of order n, $T \mapsto T^n$.
If $\mathcal{E}_n$ is a rational (resp. K3) surface, $\mathcal{E}_n/\sigma_n$ is a rational (resp. K3) surface. This follows from the comparison between the degree of the polynomials in τ (resp. T) defining $\mathcal{E}_n$ (resp. $\mathcal{E}_n/\sigma_n$).
If $\mathcal{E}_n$ has Kodaira dimension 1 it is possible that $\mathcal{E}_n/\sigma_n$ has a lower Kodaira dimension, this happens for example if $h = 2$, $k = 4$ and $n = 3$ (in this case $\mathcal{E}_n/\sigma_n$ is a K3 surface). □

The automorphisms μ_2 and ς_2 are essentially obtained by ν_2 and σ_2 with a change of coordinates on $\mathbb{P}^1$.

Proposition 5.3. *Let $\mathcal{E}$ be an elliptic fibration $y^2 = x^3 + A(\tau)x + B(\tau)$ such that*

$$A(\tau) = \tau^{4m} A(\frac{1}{\tau}) \text{ and } B(\tau) = \tau^{6m} B(\frac{1}{\tau}). \tag{4}$$

Then μ_2 and ς_2 are automorphisms of the surface,
$A(\tau) = \sum_{i=0}^{2m} a_i(t^i + t^{4m-i}) = t^{2m} \sum_{i=0}^{2m} a_i(t^{2m-i} + \frac{1}{t^{2m-i}})$ and

$B(\tau)=\sum_{j=0}^{3m}b_j(t^j+t^{6m-j})=t^{3m}\sum_{j=0}^{2m}b_j(t^{3m-j}+\frac{1}{t^{3m-j}})$.
The quotient elliptic fibrations are respectively:

$$\begin{array}{lll}\mathcal{E}/\mu_2: & Y^2= & X^3+X\sum_{i=0}^{2m}a_i\left(T^{2m-i}-\sum_{k=1}^{2m-i}\binom{2m-i}{k}T^{2m-i-2k}\right)+\\ & & +\sum_{j=0}^{3m}b_j\left(T^{3m-j}-\sum_{k=1}^{3m-j}\binom{3m-j}{k}T^{3m-j-2k}\right)\\ \mathcal{E}/\varsigma_2: & \widetilde{Y}^2= & \widetilde{X}^3+\widetilde{X}(T^2-4)^2\sum_{i=0}^{2m}a_i\left(T^{2m-i}-\sum_{k=1}^{2m-i}\binom{2m-i}{k}T^{2m-i-2k}\right)+\\ & & +(T^2-4)^3\sum_{j=0}^{3m}b_j\left(T^{3m-j}-\sum_{k=1}^{3m-j}\binom{3m-j}{k}T^{3m-j-2k}\right).\end{array}$$

Proof. Under the conditions (4) the equation of $\mathcal{E}$ is invariant under μ_2 and σ_2. The equation of the quotient surfaces are computed setting $T=t+\frac{1}{t}$, $Y=y$, $X=x$, $\widetilde{Y}=y\frac{(t+1)^3(t-1)^3}{t^6}$, $\widetilde{X}=x\frac{(t+1)^2(t-1)^2}{t^4}$. We use the equalities $t^n+\frac{1}{t^n}=T^n-\sum_{k=1}^{n}\binom{n}{k}T^{n-2k}$, $\frac{t^2\pm t+1}{t^2}=T\pm 1$, $\frac{(t\pm1)^2}{t}=T\pm 2$. □

6. AUTOMORPHISMS ON THE BASIS OF THE FIBRATION AND ELLIPTIC K3 SURFACES

In this section we consider the automorphism σ_n, Definition 5.1, acting on a K3 surface. In particular we saw (Remark 5.2) that the K3 surface X admitting σ_n as automorphism is obtained by the K3 surface $\widetilde{X/\sigma_n}$ with a base change of order n. These kind of base changes are considered for example in [Shio2], [Shio3], [Shio4], [Ku].

Remark 6.1. If $\mathcal{E}$ is an elliptic fibration on a K3 surface X admitting σ_n (resp. ς_2) as automorphism, then σ_n (resp. ς_2) is a symplectic automorphism on X. Indeed the nowhere vanishing 2-holomorphic form $dx/y\wedge d\tau$ on X is fixed by σ_n^* (resp. ς_2^*).

Proposition 6.2. *Let $2\leq n\leq 6$ and let X_n be a K3 surface which admits an elliptic fibration $\mathcal{E}_n$ satisfying the condition* (2)*. Then for a generic choice of a_i and b_i:*

- i) *X admits rank$(\Omega_{\mathbb{Z}/n\mathbb{Z}})$ independent sections of infinite order;*
- ii) *the Néron–Severi group of X_n is $NS(X_n)\simeq U\oplus\Omega_{\mathbb{Z}/n\mathbb{Z}}$;*
- iii) *the Mordell–Weil lattice of the fibration is $MW(\mathcal{E}_n)\simeq\Omega_{\mathbb{Z}/n\mathbb{Z}}$.*

Let $X_{2,2}$ be a K3 surface which admits an elliptic fibration $\mathcal{E}_{2,2}$ satisfying the conditions $A(\tau)=A(-\tau)=\tau^8A(\frac{1}{\tau})$, $B(\tau)=B(-\tau)=\tau^{12}B(\frac{1}{\tau})$ (i.e. it satisfies the conditions (2) *of Proposition 5.1 with $n=2$ and* (4) *of Proposition 5.3).*
Then for a generic choice of a_i and b_i:

- i') *$X_{2,2}$ admits 12 independent sections of infinite order;*
- ii') *the Néron–Severi group of $X_{2,2}$ is $NS(X_{2,2})\simeq U\oplus\Omega_{(\mathbb{Z}/2\mathbb{Z})^2}$;*
- iii') *the Mordell–Weil lattice of the fibration is $MW(\mathcal{E}_{2,2})\simeq\Omega_{(\mathbb{Z}/2\mathbb{Z})^2}$.*

Proof. Let p_n be the number of parameters in equation (2), m_n the moduli of the surface X_n and ρ_n the Picard number of X_n. We can act on the equation satisfying the condition (2) with the map $(x,y,\tau)\mapsto(\lambda^2x,\lambda^3y,\tau)$ (and then divide by λ), and we can act with the automorphisms of $\mathbb{P}^1$ fixing 0 and ∞ (which are the points fixed by π_n). So we have a two dimensional family of automorphisms acting on the equation of the fibrations, hence $m_n=p_n-2$. For each n we have:

(5)

n	2	3	4	5	6
m_n	10	6	4	2	2
$rk(\Omega_{\mathbb{Z}/n\mathbb{Z}})$	8	12	14	16	16

We recall that $\rho_n\leq 20-m_n$, by the construction of the moduli spaces of the polarized K3 surfaces. By Proposition 5.1 and Remark 6.1, the K3 surface X_n admits σ_n as a symplectic automorphism of order n. Hence $\Omega_{\mathbb{Z}/n\mathbb{Z}}$ is primitively embedded in $NS(X_n)$ (cf. [GS2]). The automorphism σ_n fixes the class of the zero section and the class of the fiber. These two classes span a copy of the lattice U in $NS(X_n)$, so $U\hookrightarrow NS(X_n)^{\sigma_n}$. Since $\Omega_{\mathbb{Z}/n\mathbb{Z}}=(NS(X_n)^{\sigma_n})^{\perp}$, $U\oplus\Omega_{\mathbb{Z}/n\mathbb{Z}}\hookrightarrow NS(X_n)$, in particular $\rho_X\geq 2+rk\Omega_{\mathbb{Z}/n\mathbb{Z}}$. The rank of $U\oplus\Omega_{\mathbb{Z}/n\mathbb{Z}}$ is equal to $20-m_n$ for each $n=2,\ldots,6$ (cf. table (5)), hence $\rho_X=20-m_n$ and the inclusion $U\oplus\Omega_{\mathbb{Z}/n\mathbb{Z}}\hookrightarrow NS(X_n)$ has a finite index. Since $\Omega_{\mathbb{Z}/n\mathbb{Z}}$ is primitively embedded in $NS(X_n)$ and U is a unimodular lattice, we have $NS(X_n) \simeq U \oplus \Omega_{\mathbb{Z}/n\mathbb{Z}}$. For a generic choice of the parameters a_i and b_i in the equation of X_n, the elliptic fibration has no reducible fibers (if $n \neq 5$ the singular fibers are $24I_1$, if $n = 5$ the singular fibers are $2II + 20I_1$), hence $Tr_{\mathcal{E}_n} \simeq U$ and there are no torsion sections. In particular the Mordell–Weil lattice of $\mathcal{E}_n$ is isometric to the sublattice $\phi(E(K)) \subset NS(X_n)$ (where ϕ is defined in Lemma 1.3) and $\phi(E(K)) \simeq U \oplus \Omega_{\mathbb{Z}/n\mathbb{Z}}/U \simeq \Omega_{\mathbb{Z}/n\mathbb{Z}}$. More explicitly, if F is the class of the fiber and s the class of the zero section, for each $r \in E(K)$ the map $\phi : E(K) \to NS(X) \otimes \mathbb{Q}$ sends r to $r + (2-k)F - s \in \Omega_{\mathbb{Z}/n\mathbb{Z}} \subset NS(X_n)$, where $r \cdot s = k$. The height pairing computed on $\phi(E(K))$ coincides with the intersection form on $NS(X_n)$ restricted to $\Omega_{\mathbb{Z}/n\mathbb{Z}}$. Hence the Mordell–Weil lattice is isometric to $\Omega_{\mathbb{Z}/n\mathbb{Z}}$. In particular this implies that there are $\mathrm{rank}\Omega_{\mathbb{Z}/n\mathbb{Z}}$ independent sections of infinite order. The proof in case $X_{2,2}$ is exactly the same. □

Let $2 \leq n \leq 6$ and let Y_n be the desingularization of the quotient X_n/σ_n. Then X_n is obtained by a base change of degree n of Y_n. In the following table we give the singular fibers of Y_n in the points $\tau = 0$ and $\tau = \infty$ and we describe the fibers on $\tau = 0$, $\tau = \infty$ of X_n (with $(X_n)_{\overline{\tau}}$ we denote the fiber of the fibration on X_n over the point $\overline{\tau}$.)

n	2	3	4	5	6
$(Y_n)_0, (Y_n)_\infty$	I_0^*	IV^*	III^*	II^*	II^*
$(X_n)_0, (X_n)_\infty$	$y^2 = x^3 + ax + b$	$y^2 = x^3 + 1$	$y^2 = x^3 + x$	II	II

Since $\tau = 0$ and $\tau = \infty$ are fixed points of the map $\tau \mapsto \zeta_n \tau$, σ_n acts as an automorphism of the fibers $(X_n)_0$ and $(X_n)_\infty$. Indeed for $n = 3, 4$ $(X_n)_0$ and $(X_n)_\infty$ are exactly the elliptic curves admitting an extra automorphism. The group law of the fiber of type II corresponds to the group $\mathbb{C}$.

In [Shio2], [Shio3], [Shio4] certain K3 surfaces $F^{(n)}(\alpha, \beta)$, $n = 1, \ldots, 6$ are analyzed. These are K3 surfaces admitting an elliptic fibration with equation $y^2 = x^3 - 3\alpha x + (t^n + 1/t^n - 2\beta)$. It is clear that $F^{(n)}(\alpha, \beta)$ is obtained by a base change of order n applied to $F^{(1)}(\alpha, \beta)$. As in Remark 5.2 this base change is related to the automorphism σ_n, and in fact $F^{(1)}(\alpha, \beta)$ is the quotient of $F^{(n)}(\alpha, \beta)$ by σ_n (Definition 5.1).

The surfaces described in [Shio3], [Shio4], [Shio2], [Ku] are particular members of the families described in Proposition 6.2.

Let us consider the equation $\mathcal{E}_n$ given in (3) with $n = 5$ (resp. $n = 6$), $h = 0$, $k = 1$. This is the equation of the family of K3 surface X_n in Proposition 6.2. Up to projective transformations, the equation $\mathcal{E}_5$ (resp. $\mathcal{E}_6$) becomes:

$$(6) \quad \mathcal{E}_5 : \quad y^2 = x^3 + a\tau^4 x + (\tau^{11} + b\tau^6 + \tau^1) \ \ (\text{resp. } \mathcal{E}_6 : \quad y^2 = x^3 + a'\tau^4 x + (\tau^{12} + b'\tau^6 + 1))$$

Remark 6.3. The Mordell–Weil lattice of the generic element of the family $F^{(n)}(\alpha, \beta)$ is computed in [Shio4, Theorem 2.4]. For $n = 5, 6$ the family described in [Shio4] and the family described in Proposition 6.2 are the same (this is clear considering the equation (6) of $\mathcal{E}_5$, $\mathcal{E}_6$), hence the Mordell–Weil lattice $MW(F_{gen}^{(n)})$ of [Shio4, Theorem 2.4] is the lattice $\Omega_{\mathbb{Z}/n\mathbb{Z}}$, for $n = 5, 6$. For $n < 5$ the two dimensional family described in [Shio4] is a subfamily of the one described in Proposition 6.2, and so the Mordell–Weil lattices of these two families are not isometric (moreover for $n < 5$ there are reducible fibers on $F_{gen}^{(n)}$). □

In the following we will analyze some surfaces admitting a dihedral group as group of symplectic automorphisms. This is based on the following remark.

Remark 6.4. The maps σ_n and ς_2 generate the dihedral group on n elements $\mathcal{D}_n$. So if an elliptic K3 surface admits both of them it admits the group $\mathcal{D}_n$ as group of symplectic automorphism. □

In the following we give an example of elliptic K3 surfaces admitting both the automorphisms σ_n and ς_2. Moreover we give a short proof of the fact that $\Omega_{\mathbb{Z}/n\mathbb{Z}} \simeq \Omega_{\mathcal{D}_n}$ for $n = 5, 6$. The case $n = 5$ is proved in [G2]. In the Section 8 we will consider again these isometries. Another example of elliptic K3 surfaces admitting both the automorphisms ς_2 and σ_n is considered in Section 7 where $n = 4$.

The equation of the family of elliptic K3 surfaces admitting σ_n $n=5,6$ as automorphism is given in (6). By Proposition 6.2 its Néron–Severi group is isometric to $U\oplus\Omega_{\mathbb{Z}/n\mathbb{Z}}$ and its Mordell–Weil lattice is isometric to $\Omega_{\mathbb{Z}/n\mathbb{Z}}$. The elliptic fibration (6) admits automatically also the automorphism ς_2 (Proposition 5.3). Both automorphisms σ_n and ς_2 act as the identity on the class of the fiber and on the class of the zero section and σ_n acts non trivially on all the other sections. As in Proposition 6.2, this implies that $MW(\mathcal{E})\simeq\Omega_{\mathcal{D}_n}$, but, by Proposition 6.2, $MW(\mathcal{E})\simeq\Omega_{\mathbb{Z}/n\mathbb{Z}}$ and then we conclude that $\Omega_{\mathbb{Z}/n\mathbb{Z}}\simeq\Omega_{\mathcal{D}_n}$ for $n=5,6$.

7. The group $\mathcal{D}_4$ as group of symplectic automorphisms.

Here we give an example of a family of K3 surfaces admitting the dihedral group on four elements, $\mathcal{D}_4$, as group of symplectic automorphisms. These automorphisms are induced on elliptic K3 surfaces by the automorphisms σ_4, ς_2 (Definition 5.1) hence they are related to a base change of the fibration (Remark 5.2).

In [Kl] the same family of elliptic K3 surfaces is considered in order to construct an elliptic K3 surface with 15 independent sections of infinite order. Its construction is based on two base changes (one of order 2 and one of order 4). We prove that the family of elliptic K3 admitting $\mathcal{D}_4$ as group of symplectic automorphisms that we consider is the same as the one described in [Kl]. As in Proposition 6.2, using the fact that these surfaces admit a certain group of symplectic automorphisms, we give a description of the Néron–Severi groups and of the Mordell–Weil lattices.

Remark 7.1. The elliptic K3 surfaces invariant under σ_4 and ς_2 have equations:

$$\mathcal{E}:\quad y^2=x^3+x(a\tau^8+b\tau^4+a)+(c\tau^{10}+d\tau^6+c\tau^2). \tag{7}$$

Such an equation depends on 4 parameters, but we can use the transformation $(x,y)\mapsto(\lambda^2x,\lambda^3y)$ to put one of this parameters equal to 1. Hence this family has are 3 moduli. So the Picard number of the generic K3 surface with such an elliptic fibration is at most 17. By Remark 6.4 each member of this family admits $\mathcal{D}_4$ as group of symplectic automorphisms.

Let $G=\langle\sigma_4,\varsigma_2\rangle$. Then the elliptic fibration $\mathcal{E}/G$ has equation

$$\mathcal{E}/G:\quad Y^2=X^3+X(T^2-4)^2(aT+b)+(T^2-4)^3(cT+d),$$

where $T=(t^4+1/t^4)$, $X=x(t^2+1/t^6)$, $Y=y(t-1/t^7)$.

For a generic choice of a,b,c,d the elliptic fibration $\mathcal{E}/G$ has $III^*+2I_0^*+3I_1$ as singular fibers and does not admit sections of infinite order. Indeed the trivial lattice of this fibration has rank 17 and thus the Picard number is at least 17. Since the Picard number of two K3 surfaces are the same if there exists a dominant rational map between them, the Picard number of this surface and of the one with fibration $\mathcal{E}$ is exactly 17. $\square$

Remark 7.2. The elliptic fibration $\mathcal{E}/G$ has the properties of the elliptic fibration $\widetilde{\pi}:\widetilde{X}\rightarrow\mathbb{P}^1$ considered in [Kl, Table 2]. In particular these elliptic fibrations coincide under the condition given in [Kl, Proposition 3.2] that the rank of a certain twist (in [Kl, Construction 3.1]) is 0. This implies that the elliptic fibration $\mathcal{E}$ is the one described in [Kl, Theorem 1.1] as elliptic K3 with Mordell–Weil rank 15. Indeed this fibration is constructed by Kloosterman performing two base changes on $\mathcal{E}/G$ (one of order 4 and one of order 2), which correspond to the presence of the symplectic automorphisms σ_4 and ς_2 on $\mathcal{E}$. $\square$

In the following we construct an elliptic fibration (which is a special member of the family $\mathcal{E}$ given in Remark 7.1) admitting $\mathcal{D}_4$ as group of symplectic automorphisms and use it to determine the lattice $\Omega_{\mathcal{D}_4}$. We will use the description of this lattice to find the Mordell–Weil lattice and the Néron–Severi group of the surface constructed in [Kl].

7.1. **Kummer surface of the product of two elliptic curves.** Let us recall the construction of a Kummer surface $Km(A)$. Let A be an Abelian surface. Let $\iota_A:a\mapsto -a$. Hence ι is an involution fixing the sixteen point of $A[2]=\{a\in A \text{ such that } 2a=0\}$. The quotient A/ι is a singular surface with sixteen singularities of type A_1. Let us denote with $Km(A)$ the desingularization of this surface.

The surface $Km(A)$ is a K3 surface, called Kummer surface of A. We have the following commutative diagram:

$$\begin{tikzcd}
\iota \circlearrowleft A \arrow[d, "\pi"'] & \widetilde{A} \circlearrowleft \widetilde{\iota} \arrow[l, "\beta"'] \arrow[d, "\widetilde{\pi}"] \\
A/\iota & Km(A) \simeq \widetilde{A}/\widetilde{\iota} \arrow[l]
\end{tikzcd}$$

β is the blow up of A in the points of $A[2]$, $\widetilde{\iota}$ is the involution induced by ι on $\widetilde{A}$, π and $\widetilde{\pi}$ are the quotient maps.

Let C and D two elliptic curves with equation:

$$\begin{aligned} C:&\ \ v^2 = u^3 + Au + B \\ D:&\ \ \nu^2 = f(\tau)\ \ deg(f(\tau)) = 4, 3. \end{aligned} \tag{8}$$

Then (as in [Ku]) we consider the equation $f(\tau)y^2 = x^3 + Ax + B$, or, which is the same under the transformation $y \mapsto y/f^2(\tau)$ and $x \mapsto x/f^3(\tau)$, the equation

$$y^2 = x^3 + Af^2(\tau)x + Bf^3(\tau). \tag{9}$$

Let $f : \mathcal{E} \to \mathbb{P}^1$ be the elliptic fibration with equation (9).
The fibration $\mathcal{E}$ is an isotrivial fibration with fibers isomorphic to C. Indeed under the change of coordinates

$$\left\{ \begin{array}{l} y = \nu^3 v \\ x = \nu^2 u. \end{array} \right. \tag{10}$$

the equation (9) becomes $\nu^6 v^2 = \nu^6 u^3 + Af^2(\tau)\nu^2 u + Bf^3(\tau)$. Since $\nu^2 = f(\tau)$, we obtain that the fiber over each fixed $\overline{\tau}$ has equation $\nu^6(v^2 = u^3 + Au + B)$.

The equation (9) is the equation of an elliptic fibration on the Kummer surface $Km(C \times D)$. In fact let us consider the trivial elliptic fibration $C \times D \to D$ defined over the Abelian surface $C \times D$ and the map $\varphi : D \to \mathbb{P}^1$ defined as $(\tau, \nu) \mapsto (\tau, -\nu)$. Then we have the following commutative diagram:

$$\begin{tikzcd}
\Phi: & C \times D \arrow[r, "2:1"] \arrow[d] & \mathcal{E} \arrow[d, "f"] \\
\varphi: & D \arrow[r, "2:1"] & \mathbb{P}^1_\tau
\end{tikzcd}$$

where the map Φ is induced by the map φ and hence it has degree 2. The explicit equation of Φ descends by (10) and is

$$\Phi : ((u, v), (\nu, \tau)) \mapsto (x, y, \tau) = (\nu^2 u, \nu^3 v, \tau).$$

The points $(p, q) = ((u, v), (\tau, \nu))$, $(p', q') = ((a, b), (\alpha, \beta)) \in C \times D$ are such that $\Phi(p, q) = \Phi(p', q')$ if and only if either $p = p'$ and $q = q'$, or $u = a$, $v = -b$, $\tau = \alpha$, $\nu = -\beta$. The maps $\iota_C : c \mapsto -c$, $\iota_D : d \mapsto -d$ is represented in coordinates by

$$\left\{ \begin{array}{lll} (u, v) \mapsto (u, -v) & \text{on the elliptic curve} & C \\ (\tau, \nu) \mapsto (\tau, -\nu) & \text{on the elliptic curve} & D \end{array} \right.$$

Hence the map Φ identifies the point (p, q) with $(-p, -q)$, so it coincides with the map $\iota_{C\times D} : a \mapsto -a$ on the Abelian surface $C \times D$. In particular, the desingularization of the quotient $C \times D/\iota_{C\times D}$ is the surface $Km(C \times D)$ and the singular elliptic fibration $\mathcal{E}$ induces an elliptic fibration over $Km(C \times D)$. With an abuse of notation we will call $\mathcal{E}$ also the elliptic fibration over $Km(C \times D)$.

7.2. $Km(E_i \times E_i)$**.** From now on we assume that the two elliptic curves C and D are E_i, i.e. they admit an automorphism of order 4 which is not a translation.

Remark 7.3. Since the elliptic curves $C \simeq D$ are isomorphic to E_i, the equations (8) can be chosen on the following way

$$\begin{aligned} C:&\ \ v^2 = u^3 + u \\ D:&\ \ \nu^2 = \tau^4 - 1 \end{aligned}$$

where the automorphisms of order 4 are $\varphi_C : (u,v) \mapsto (-u, iv)$ and $\varphi_D : (\tau,\nu) \mapsto (i\tau,\nu)$ respectively. Hence the equation (9) becomes

$$y^2 = x^3 + (\tau^4 - 1)^2 x. \tag{11}$$

The zero of the group law on the curve C (resp. D) has to be fixed by the map ι_C (resp. ι_D) hence we can assume that it is the point at infinity (resp. the point $(\tau,\nu) = (1,0)$). $\square$

We observe that this elliptic fibration is a special member of the family described in Remark 7.1, indeed its equation is obtained by equation (7) putting $a = 1$, $b = -2$, $c = d = 0$. Hence it admits the automorphisms σ_4 and ς_2 and so the group $\mathcal{D}_4$ is a group of symplectic automorphisms on it. We now relate the automorphisms σ_4 and ς_2 with automorphisms of order 4 on the elliptic curves C and D.

The points of $C[2]$ are, in homogeneous coordinates, $0_C = (0:1:0)$, $c_1 = (0:0:1)$, $c_2 = (i:0:1)$, $c_3 = (-1:0:1)$. They correspond to the image of the point $0_C = 0$, $c_1 = (1+i)/2$, $c_2 = 1/2$, $c_3 = i/2$ under the map $\eta : \mathbb{C} \rightarrow \mathbb{C}/\mathbb{Z}[i] = E_i$. They can be identified with the points $0_C = (0,0)$, $c_1 = (1,1)$, $c_2 = (1,0)$, $c_3 = (0,1)$ under the identification of $C[2]$ with $(\mathbb{Z}/2\mathbb{Z})^2$.
The points of $D[2]$ are the points fixed by the map ι_D. Hence they are $0_D = (1;0)$, $d_1 = (i;0)$, $d_2 = (-1;0)$, $d_3 = (-i;0)$. They are the image of the points $0_D = (0,0)$, $d_1 = i/2$, $d_2 = (1+i)/2$, $d_3 = 1/2$ under η and they can be identified with the points $0_D = (0,0)$, $d_1 = (0,1)$, $d_2 = (1,0)$, $d_3 = (1,1)$ under the identification of $D[2]$ with $(\mathbb{Z}/2\mathbb{Z})^2$.
The map φ_C fixes the points 0_C and c_1 and switches the point c_2 and c_3. On $\mathbb{C}$ it corresponds to $z_1 \mapsto iz_1$.
The map φ_D acts in the following way: $0_D \mapsto d_1 \mapsto d_2 \mapsto d_3$. On $\mathbb{C}$ it corresponds to $z_2 \mapsto iz_2 + 1/2$.

We will denote with $K_{e,f,g,h}$ the rational curve in $Km(E_i \times E_i)$ which is the exceptional curve over the points $p_{e,f,g,h} = (e,f) \times (g,h) \in (E_i \times E_i)[2]$.
Let $\Delta = \{(c,d) \in Km(E_i \times E_i) | c = d\}$ and $\Gamma = \{(c,d) \in Km(E_i \times E_i) | c = \varphi_{E_i}(d)\}$ where φ_{E_i} is the automorphism of order four of E_i. They correspond to two classes in Néron–Severi group.

Remark 7.4. We identify C with $\mathbb{R}^2_{x_1,x_2}/\mathbb{Z}^2$ (i.e. the real coordinates of $\mathbb{R}^2$ will be denoted by x_1 and x_2), and D with $\mathbb{R}^2_{x_3,x_4}/\mathbb{Z}^2$. Since $dx_i \wedge dx_j$, $i<j$ is a basis of $H^2(C \times D, \mathbb{Z})$, we can express the classes $[C]$, $[D]$, $[\Gamma]$, $[\Delta]$ as linear combination of $dx_i \wedge dx_j$. More precisely

$$\begin{array}{rclrcl} [C] & = & dx_3 \wedge dx_4, & [\Gamma] & = & dx_1 \wedge dx_2 + dx_3 \wedge dx_4 + dx_1 \wedge dx_3 + dx_2 \wedge dx_4, \\ [D] & = & dx_1 \wedge dx_2, & [\Delta] & = & dx_1 \wedge dx_2 + dx_3 \wedge dx_4 + dx_2 \wedge dx_3 - dx_1 \wedge dx_4. \end{array}$$

This computation is done as in [G1, Remark 3.11]. $\square$

Remark 7.5. Let $\omega_{i,j} = \pi_*(\beta^*(dx_i \wedge dx_j))$. By Remark 7.4, we have

$$\begin{array}{rclrcl} \pi_*(\beta^*[C]) & = & \omega_{3,4}, & \pi_*(\beta^*[\Gamma]) & = & \omega_{1,2} + \omega_{3,4} + \omega_{1,3} + \omega_{2,4}, \\ \pi_*(\beta^*[D]) & = & \omega_{1,2}, & \pi_*(\beta^*[\Delta]) & = & \omega_{1,2} + \omega_{2,3} + \omega_{3,4} - \omega_{1,4}. \end{array}$$

Hence:

$$\begin{array}{rcl} \widetilde{C_0} := \pi(\widetilde{C \times (0,0)}) & = & \frac{1}{2}\left(\omega_{3,4} - \sum_{(a,b)\in(\mathbb{Z}/2\mathbb{Z})^2} K_{a,b,0,0}\right), \\ \widetilde{D_0} := \pi(\widetilde{(0,0) \times D}) & = & \frac{1}{2}\left(\omega_{1,2} - \sum_{(a,b)\in(\mathbb{Z}/2\mathbb{Z})^2} K_{0,0,a,b}\right), \\ \widetilde{\Gamma} := \pi(\widetilde{\Gamma}) & = & \frac{1}{2}\left(\omega_{1,2} + \omega_{3,4} + \omega_{1,3} + \omega_{2,4} - \sum_{(a,b)\in(\mathbb{Z}/2\mathbb{Z})^2} K_{a,b,b,a}\right), \\ \widetilde{\Delta} := \pi(\widetilde{\Delta}) & = & \frac{1}{2}\left(\omega_{1,2} + \omega_{2,3} + \omega_{3,4} - \omega_{1,4} - \sum_{(a,b)\in(\mathbb{Z}/2\mathbb{Z})^2} K_{a,b,a,b}\right). \end{array}$$

$\square$

We use the following convention: if W is a vector subspace of the affine space $(C \times D)[2] = (\mathbb{Z}/2\mathbb{Z})^4$, then $\bar{K}_W$ is the class $\frac{1}{2}\sum_{a\in W} K_a$ and $\hat{K} = \frac{1}{2}\sum_{a\in(\mathbb{Z}/2\mathbb{Z})^4} K_a = \bar{K}_{A[2]}$. Moreover, $W_i = \{a = (a_1,a_2,a_3,a_4) \in A[2]$ such that $a_i = 0\}$, $i = 1,2,3,4$.

Proposition 7.6. *Let $Km(C\times D)$ be the K3 surface admitting the elliptic fibration*

$$y^2=x^3+(\tau^4-1)^2x.$$

A $\mathbb{Z}$-basis for $NS(Km(C\times D))$ is given by the classes:

$$\begin{array}{cccccccccc} K_{0000}, & K_{0001}, & K_{0010}, & K_{0100}, & K_{1000}, & K_{0011}, & K_{0101}, & K_{1001}, & K_{0110}, & K_{1010},\\ K_{1100}, & \bar{K}_{W_4}, & \bar{K}_{W_3}, & \bar{K}_{W_2}, & \bar{K}_{W_1}, & \hat{K}, & \widetilde{C_0}, & \widetilde{D_0}, & \widetilde{\Delta}, & \widetilde{\Gamma}. \end{array}$$

A $\mathbb{Z}$-basis for $T_{Km(C\times D)}$ is $\omega_{1,4}+\omega_{2,3}$, $\omega_{1,3}-\omega_{2,4}$. and $T_{Km(C\times D)}\simeq\langle 4\rangle^2$.
Let $\mathcal{D}_4=\langle\sigma_4,\varsigma_2\rangle$. The lattice $NS(Km(C\times D))^{\mathcal{D}_4}$ is generated by the classes

$$\bar{K}_{W_1}+\bar{K}_{W_2},\quad \omega_{3,4},\quad \widetilde{D_0},\quad K_{0000}+K_{0001}+K_{0010}+K_{0011},\quad \hat{K}.$$

The lattice $\Omega_{\mathcal{D}_4}:=(H^2(Km(C\times D))^{\mathcal{D}_4})^{\perp}=(NS(Km(C\times D))^{\mathcal{D}_4})^{\perp}$ is generated by the classes

$$\begin{array}{ccc} \widetilde{\Gamma}-2K_{0010}-\widetilde{C_0}-\widetilde{D_0}, & \widetilde{\Delta}-2K_{0010}-\widetilde{C_0}-\widetilde{D_0}, & K_{0001}-K_{0010},\\ K_{0000}-K_{0010}, & \hat{K}-2K_{0010}-4K_{0100}-2K_{1100}, & \bar{K}_{W_1}-2K_{0010}-2K_{0100},\\ \bar{K}_{W_2}-2K_{0010}-2K_{0100}, & \bar{K}_{W_3}-K_{0010}-2K_{0100}-K_{1100}, & \bar{K}_{W_4}-K_{0010}-2K_{0100}-K_{1100},\\ K_{1000}-K_{0100}, & K_{1010}-K_{0100}, & K_{0110}-K_{0100},\\ K_{1001}-K_{0100}, & K_{0101}-K_{0100}, & K_{0011}-K_{0010}. \end{array}$$

The lattice $\Omega_{\mathcal{D}_4}$ has rank 15, its discriminant is -1024 and its discriminant group is $(\mathbb{Z}/4\mathbb{Z})^5$.
The lattice $H^2(Km(C\times D),\mathbb{Z})^{\mathcal{D}_4}$ is isometric to $NS(Km(C\times D))^{\mathcal{D}_4}\oplus T_{Km(C\times D)}$.

Proof. The results on $NS(Km(C\times D))$ and $T_{Km(C\times D)}$ are trivial consequences of Remark 7.5, of the description of the Kummer lattice given, for example, in [BPV, Chapter VIII, Section 5], [PS, Appendix 5] and more in general of the known description of the classes generating the second cohomology group of a Kummer surface (see for example, [G1]).
A direct computation on the coordinates (x,y,τ) shows that the automorphism $\varphi_C^3\times\varphi_D$ induces the automorphism σ_4 on $Km(C\times D)$. The automorphism $\varphi_C^3\times\varphi_D$ acts on $C\times D$ as $(z_1,z_2)\mapsto(-iz_1,iz_2+1)$. So the action of σ_4 on the curves of the lattice K (induced by the one of $\varphi_C^2\times\varphi_D$ on $(C\times D)[2]$) is

$$\begin{array}{c||c} K_{0,0,0,0}\rightarrow K_{0,0,0,1}\rightarrow K_{0,0,1,1}\rightarrow K_{0,0,1,0} & K_{0,1,0,0}\rightarrow K_{1,0,0,1}\rightarrow K_{0,1,1,1}\rightarrow K_{1,0,1,0}\\ K_{1,0,0,0}\rightarrow K_{0,1,0,1}\rightarrow K_{1,0,1,1}\rightarrow K_{0,1,1,0} & K_{1,1,0,0}\rightarrow K_{1,1,0,1}\rightarrow K_{1,1,1,1}\rightarrow K_{1,1,1,0}. \end{array}$$

Since

$$\sigma_4(\omega_{1,2})=\omega_{1,2},\ \sigma_4(\omega_{1,3})=-\omega_{2,4},\ \sigma_4(\omega_{1,4})=\omega_{2,3},\ \sigma_4(\omega_{2,3})=\omega_{1,4},\ \sigma_4(\omega_{2,4})=-\omega_{1,3},\ \sigma_4(\omega_{3,4})=\omega_{3,4},$$

the classes $\pi_*(\beta^*[C])$, $\pi_*(\beta^*[D])$ are fixed and

$$\begin{array}{rl} \sigma_4\left(\pi_*(\beta^*[\Delta])\right)= & -\pi_*(\beta^*[\Delta])+2\pi_*(\beta^*[C])+2\pi_*(\beta^*[D]),\\ \sigma_4\left(\pi_*(\beta^*[\Gamma])\right)= & -\pi_*(\beta^*[\Gamma])+2\pi_*(\beta^*[C])+2\pi_*(\beta^*[D]). \end{array}$$

Let ψ_D be the automorphism, of the curve D, $\psi_D:(\tau,\nu)\rightarrow(1/\tau,i\nu/\tau^2)$. The automorphism $\varphi_C^3\times\psi_D$ induces on $Km(C\times D)$ the automorphism ς_2. We observe that φ_C^3 and ψ_D have order four respectively on C and D. However $\varphi_C^3\times\psi_D$ induces an automorphisms of order two on $Km(C\times D)$, indeed $(\varphi_C^3\times\psi_D)^2=\iota_C\times\iota_D$ which is the identity on $Km(C\times D)$.
The automorphism ψ_D of D has order four and fixes the points $0_D=(1,0)$ and $(-1,0)$. Hence it corresponds to $z_2\mapsto iz_2$. So the automorphism ς_2 is induced by the automorphism $(z_1,z_2)\mapsto(-iz_1,iz_2)$ on $C\times D$. The action of ς_2 on the curves of the Kummer lattice is

$$\begin{array}{lllllllll} K_{0,0,0,1} & \leftrightarrow & K_{0,0,1,0}; & K_{0,1,0,0} & \leftrightarrow & K_{1,0,0,0}; & K_{0,1,0,1} & \leftrightarrow & K_{1,0,1,0};\\ K_{0,1,1,0} & \leftrightarrow & K_{1,0,0,1}; & K_{0,1,1,1} & \leftrightarrow & K_{1,0,1,1}; & K_{1,1,0,1} & \leftrightarrow & K_{1,1,1,0} \end{array}$$

and the other classes of the Kummer lattice are fixed. The action of ς_2 on the classes $\pi_*(\beta^*[C])$, $\pi_*(\beta^*[D])$, $\pi_*(\beta^*[\Delta])$, $\pi_*(\beta^*[\Gamma])$ is the same as the action of σ_4.
Since we described the action of σ_4 and ς_2, it is trivial to compute $\Omega_{\mathcal{D}_4}$ and its orthogonal in $H^2(Km(C\times D),\mathbb{Z})$. □

Proposition 7.7. *Let X be the generic elliptic K3 surface admitting the automorphisms σ_4 and ς_2, i.e. X is the generic K3 surface with equation* (7). *Then $NS(X) \simeq U \oplus \Omega_{\mathcal{D}_4}$ and $MW(X) \simeq \Omega_{\mathcal{D}_4}$.*

Proof. In [W] is proved that the action of a finite group of symplectic automorphisms on the second cohomology group of the K3 surface (and hence on Λ_{K3}) is essentially unique (cf. [N1, Definition 4.6]). Since the action of G on $H^2(X,\mathbb{Z})$ does not depend on X up to isometry, the lattice $(H^2(X,\mathbb{Z})^G)^{\perp} =: \Omega_G$ does not depend on X. In particular this implies that if G is a finite group of symplectic automorphisms of a K3 surface S, the lattice Ω_G is primitively embedded in $NS(S)$, indeed since G is symplectic $T_S \hookrightarrow H^2(S,\mathbb{Z})^G$ and hence $NS(S) \hookleftarrow (H^2(S,\mathbb{Z})^G)^{\perp}$. In Proposition 7.6 we computed the lattice $\Omega_{\mathcal{D}_4}$. It follows that $\Omega_{\mathcal{D}_4} \hookrightarrow NS(X)$. The proposition follows as in the proof of Proposition 6.2. □

Remark 7.8. Proposition 7.7 gives the Néron–Severi group and the Mordell–Weil lattice of the elliptic K3 surface admitting 15 independent sections described in [Kl, Theorem 1.1] (cf. Reamrk 7.2). □

The paper [GL] classifies lattices which have dihedral groups $\mathcal{D}_n$ (for $n = 2,3,4,5,6$) in the group of the isometries and which have the properties:

- the lattices are rootless (i.e. there are no elements of length 2),
- they are the sum of two copies of $E_8(2)$,
- there are involutions in $\mathcal{D}_n$ acting as minus the identity on each copy of $E_8(2)$.

In [GL, Section 5.2] it is proved that there are three lattices satisfying these properties and admitting $\mathcal{D}_4$ as group of isometries, they are called $DIH_8(15)$, $DIH_8(16, DD4)$, $DIH_8(16,0)$.

Remark 7.9. The lattice $DIH_8(15)$ described in [GL] is isometric to the lattice $\Omega_{\mathcal{D}_4}(-1)$.
In fact $\Omega_{\mathcal{D}_4}$ has no vectors of length -2 (this is a property of all the lattices Ω_G, cf. [N1, Lemma 4.2], [W]). The two involutions ς_2 and $\varsigma_2\sigma_4$ generate the group $\mathcal{D}_4$ as group of isometries on $\Omega_{\mathcal{D}_4}$. Moreover they act as minus identity on two copies of $E_8(-2)$ (because they are both symplectic involutions, so they act as minus the identity on $\Omega_{\mathbb{Z}/2\mathbb{Z}}$ and $\Omega_{\mathbb{Z}/2\mathbb{Z}} \simeq E_8(-2)$, cf. [vGS, Section 1.3]) and the sum of these two copies of $E_8(-2)$ is $\Omega_{\mathcal{D}_4}$ (this follows from the explicit computation on the action of ς_2 and $\varsigma_2\sigma_4$ on $\Omega_{\mathcal{D}_4}$), hence $\Omega_{\mathcal{D}_4}(-1)$ is isometric to one of the lattices $DIH_8(15)$, $DIH_8(16, DD4)$, $DIH_8(16,0)$. Since $\Omega_{\mathcal{D}_4}$ has rank 15 and the lattices $DIH_8(15)$, $DIH_8(16, DD4)$, $DIH_8(16,0)$ have rank 15, 16, 16 respectively, $\Omega_{\mathcal{D}_4}(-1) \simeq DIH_8(15)$. □

8. Dihedral groups of automorphisms on elliptic fibrations with torsion.

In [G2] we proved that if a K3 surface admits $G = \mathbb{Z}/5\mathbb{Z}$ as group of symplectic automorphisms, then the same surface admits $\mathcal{D}_5$ as group of symplectic automorphisms. Moreover it is proved that $\Omega_{\mathbb{Z}/5\mathbb{Z}}$ is isometric to $\Omega_{\mathcal{D}_5}$. A similar phenomenon happens also for other groups of symplectic automorphisms. Here we will consider these other cases and we reconsider the situation described in [G2].
We consider the following groups with their presentation:

$$\begin{array}{c}\mathcal{D}_5 = \langle c,d \mid c^5 = 1,\ d^2 = 1, cd = dc^{-1}\rangle,\\ \mathcal{D}_6 = \langle c,d \mid c^6 = 1,\ d^2 = 1, cd = dc^{-1}\rangle\\ \mathbb{Z}/2\mathbb{Z} \times \mathcal{D}_4 = \langle c,d,e \mid c^2 = 1,\ d^2 = 1, e^4 = 1,\ ce = ec,\ cd = dc,\ ed = de^{-1}\rangle\\ J := \langle c,d,e \mid c^2 = 1,\ d^3 = 1, e^3 = 1,\ cd = d^{-1}c,\ ce = e^{-1}c\rangle.\end{array}$$

Let G be one of the following four groups: $\mathbb{Z}/n\mathbb{Z}$, $n = 5,6$, $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$, $(\mathbb{Z}/3\mathbb{Z})^2$. In [GS2] the equation of the generic elliptic K3 surface X_G admitting G as torsion part of Mordell–Weil group is given. For these groups the equation of elliptic fibration $\mathcal{E}_G$ on X_G depends on the choice of two polynomials of degree 2. More precisely the equation of $\mathcal{E}_G$ is of type

$$y^2 = x^3 + C(a,b)x^2 + A(a,b)x + B(a,b)$$

where $a(\tau) = \widetilde{a}(\tau,1)$, $b(\tau) = \widetilde{b}(\tau,1)$, $\widetilde{a}(\tau,\sigma)$, $\widetilde{b}(\tau,\sigma)$ are two homogenous polynomials of degree two in τ, σ, and A, B, C are homogenous polynomials in a, b of degree 4,6,2 respectively (cf. [GS2, Tables 1 and 2]).

Proposition 8.1. *Let X_G be the K3 surface admitting the generic elliptic fibration $\mathcal{E}_G$ such that $MW(\mathcal{E}_G)=G$. Let G be one of the four groups $\mathbb{Z}/n\mathbb{Z}$, $n=5,6$, $\mathbb{Z}/2\mathbb{Z}\times\mathbb{Z}/4\mathbb{Z}$, $(\mathbb{Z}/3\mathbb{Z})^2$ and let t_1 be the a generator of $\mathbb{Z}/n\mathbb{Z}$, $n=5,6$, of $\mathbb{Z}/4\mathbb{Z}$, of a copy of $\mathbb{Z}/3\mathbb{Z}$ respectively. Then there exists a symplectic involution ι on X_G and this involution is such that $\mathcal{D}_m=\langle\sigma_{t_1},\iota\rangle$ where m is the order of t_1.*
Moreover

$$\Omega_{\mathbb{Z}/n\mathbb{Z}}\simeq\Omega_{\Omega_n} \text{ for } n=5,6, \quad \Omega_{\mathbb{Z}/2\mathbb{Z}\times\mathbb{Z}/4\mathbb{Z}}\simeq\Omega_{\mathbb{Z}/2\mathbb{Z}\times\mathcal{D}_4}, \quad \Omega_{\mathbb{Z}/3\mathbb{Z}\times\mathbb{Z}/3\mathbb{Z}}\simeq\Omega_J. \tag{12}$$

Proof. First we will prove that the elliptic fibration $\mathcal{E}_G$ admits a symplectic involution. To construct this involution, we consider an automorphism on $\mathbb{P}^1$.
Step 1: construction of an involution on $\mathbb{P}^1$. Let $\widetilde{a}(\tau,\sigma)$, $\widetilde{b}(\tau,\sigma)$ be two homogeneous polynomials of degree two. Let α_1 and α_2 be the roots of $\widetilde{a}$ and β_1, β_2 the ones of $\widetilde{b}$. There is an involution θ of $\mathbb{P}^1$ such that $\theta(\alpha_1)=\alpha_2$ and $\theta(\beta_1)=\beta_2$. Indeed up to an isomorphism of $\mathbb{P}^1$ one can always suppose that $\widetilde{a}(\tau,\sigma)=\tau\sigma$ and $\widetilde{b}(\tau,\sigma)=\lambda(\tau-\sigma)(\tau-\mu\sigma)$. Hence the involution $\theta(\tau,\sigma)=(\mu\sigma,\tau)$ has the required property. We note that on the affine coordinate τ the action of θ is $\tau\mapsto\mu/\tau$.
Step 2: construction of a symplectic involution on X_G. The automorphism

$$\vartheta:(x,y,\tau)\mapsto(\frac{\mu^2x}{\tau^4},-\frac{\mu^3y}{\tau^6},\frac{\mu}{\tau})=(\frac{\mu^2x}{\tau^4},-\frac{\mu^3y}{\tau^6},\theta(\tau))$$

is a symplectic automorphism of X_G. Indeed, $\mathcal{E}_G$ has an equation of type:

$$y^2=x^3+C(a(\tau),b(\tau))x^2+A(a(\tau),b(\tau))x+B(a(\tau),b(\tau)).$$

Since $a(\theta(\tau))=\frac{\mu}{\tau^2}a(\tau)$ and $b(\theta(\tau))=\frac{\mu}{\tau^2}b(\tau)$ it is clear that the equation is invariant under ϑ. Moreover, this automorphism is symplectic, in fact $\vartheta^*(dx/y\wedge d\tau)=dx/y\wedge d\tau$.
Step 3: description of the group $\langle\sigma_{t_1},\iota\rangle$ Let t_1 be a section of order $5,6,4,3$ if G is $\mathbb{Z}/5\mathbb{Z}$, $\mathbb{Z}/6\mathbb{Z}$, $\mathbb{Z}/2\mathbb{Z}\times\mathbb{Z}/4\mathbb{Z}$, $(\mathbb{Z}/3\mathbb{Z})^2$ respectively. Let $\iota=\sigma_{t_1}\circ\vartheta$, then ι is a symplectic automorphism on X_G, because it is composition of two symplectic automorphisms. Let F_s be the fiber over the point $s\in\mathbb{P}^1$. The fibers F_τ and $F_{\iota_{\mathbb{P}^1}(\tau)}$ are isomorphic and we chose an isomorphism such that $t_i(\tau)=t_i(\iota_{\mathbb{P}^1}(\tau))$ for each section t_i, hence to each point $P_\tau\in F_\tau$ corresponds uniquely a point $P_{\iota_{\mathbb{P}^1}(\tau)}\in F_{\iota_{\mathbb{P}^1}(\tau)}$. The automorphism ι acts in the following way $P_\tau\mapsto -P_{\iota_{\mathbb{P}^1}(\tau)}+T_{\iota_{\mathbb{P}^1}(\tau)}$ where '$-$' is the operation on the elliptic curve and T_s is the intersection of the fiber F_s with the section t_1. Its clear that ι is an involution and its action on the section is $\iota(t_i)=t_{-i+1}$. This involution, together with σ_{t_1}, generates the dihedral group $\mathcal{D}_n$ if t_1 is an n-torsion section.
Moreover, if $G=\mathbb{Z}/2\mathbb{Z}/\times\mathbb{Z}/4\mathbb{Z}$ and r is the 2-torsion section such that $r\notin\langle t_1\rangle$, then $\iota\sigma_r=\sigma_r\iota$ and hence $\langle\sigma_r,\sigma_t,\iota\rangle=\mathbb{Z}/2\mathbb{Z}\times\mathcal{D}_4$.
If $G=(\mathbb{Z}/3\mathbb{Z})$ and r is a 3-torsion section such that $r\notin\langle t_1\rangle$, then $\iota\sigma_r=\sigma_r^{-1}\iota$, hence $\langle\sigma_r,\iota\rangle\simeq\langle\sigma_t,\iota\rangle=\mathcal{D}_3$ and $\langle\sigma_r,\sigma_t,\iota\rangle\simeq J$.
Step 4: proof of (12). The automorphism ι fixes the class of the fiber on $\mathcal{E}_G$ and the sum of all the sections (it clearly sends sections to sections). Since ι does not fix the points of $\mathbb{P}^1$ corresponding to singular fibers (this follows from the computation of the discriminant of the elliptic fibration $\mathcal{E}_G$), there are no components of the reducible fibers which are fixed by ι. Hence

$$NS(X)^\iota\otimes\mathbb{Q}=\langle F,\sum_{t\in MW(\mathcal{E}_G)}t\rangle\otimes\mathbb{Q} \quad\text{and so}\quad (NS(X)^\iota)^\perp=\langle F,\sum_{t\in MW(\mathcal{E}_G)}t\rangle^\perp.$$

We recall that also $NS(X)^G\otimes\mathbb{Q}=\langle F,\sum_{t\in MW(\mathcal{E}_G)}t\rangle\otimes\mathbb{Q}$ (cf. [GS1], [GS2]) and hence

$$\Omega_G\simeq(NS(X)^G)^\perp\simeq\langle F,\sum_{t\in MW(\mathcal{E}_G)}t\rangle^\perp.$$

Let $H=\langle\iota,G\rangle$. Since $NS(X)^G\otimes\mathbb{Q}=NS(X)^\iota\otimes\mathbb{Q}$, we have $NS(X)^H\otimes\mathbb{Q}=NS(X)^G\otimes\mathbb{Q}$ and so

$$\Omega_H\simeq(NS(X)^H)^\perp\simeq(NS(X)^G)^\perp\simeq\Omega_G.$$

□

Remark 8.2. The group J is isomorphic to the group $\mathfrak{A}_{3,3}$ described in [Mu]. Indeed J is a finite group of symplectic automorphisms on a K3 surface. These groups are listed by Xiao, [X, Table 2]. The order of J is 18 and there are only two groups in the Xiao's list with order 18. One of them is $\mathbb{Z}/3\mathbb{Z} \times \mathcal{D}_3$ but in J there are no elements of order 3 commuting with an involution, hence H has to be isomorphic to the other one, which is $\mathfrak{A}_{3,3}$. $\square$

In [G2] the case of the elliptic fibration $\mathcal{E}_{\mathbb{Z}/5\mathbb{Z}}$ is analyzed. In this situation it is possible to show that $E_8(-1) \oplus E_8(-1) \subset NS(X_{\mathbb{Z}/5\mathbb{Z}})$. Hence one can define an involution which switches these two copies of $E_8(-1)$ (by Definition this is a Morrison–Nikulin involution, cf. [vGS, Section 4.4]). In the case $G = \mathbb{Z}/6\mathbb{Z}$ the situation is very similar.
We will need the following trivial lemma and so we recall that given a lattice T, the length $l(A_T)$ is is the minimum number of generator of $A_T := T^\vee/T$.

Lemma 8.3. *Let $n \in \mathbb{N}$ and (T, b_T) be a lattice such that: i) $rkT = l(A_T)$;*
ii) $T^\vee/T \simeq \bigoplus_i \mathbb{Z}/nd_i\mathbb{Z}$ and β_i generate $\mathbb{Z}/nd_i\mathbb{Z}$ in $T^\vee/T$.
Let L be the free $\mathbb{Z}$-module $L = \{x \in T \otimes \mathbb{Q}$ such that $nx \in T\}$ and $b_L = n(b_{T\otimes\mathbb{Q}})_{|L}$ a bilinear form defined on L. Then (L, b_L) is a lattice and $T \simeq L(n)$ if and only if $b_{T\otimes\mathbb{Q}}(d_i\beta_i, d_j\beta_j) \in \frac{1}{n}\mathbb{Z}$.
Moreover L is an even lattice if and only if $b_{T\otimes\mathbb{Q}}(d_i\beta_i, d_i\beta_i) \in \frac{2}{n}\mathbb{Z}$.

Proposition 8.4. *Let G be one of the following four groups: $\mathbb{Z}/n\mathbb{Z}$, $n = 5, 6$, $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$, $(\mathbb{Z}/3\mathbb{Z})^2$. Let $\mathcal{E}_G$ be the generic elliptic fibration such that $MW(\mathcal{E}_G) = G$. Then: the involution ι (defined in proof of Proposition 8.1) is a Morrison–Nikulin involution for $G = \mathbb{Z}/n\mathbb{Z}$, $n = 5, 6$ and $G = (\mathbb{Z}/3\mathbb{Z})^2$ and it is* **not** *a Morrison–Nikulin involution if $G = \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$.*

Proof. The case $G = \mathbb{Z}/n\mathbb{Z}$, $n = 5, 6$. We consider only the case $n = 6$ because the other is very similar and in part it is analyzed in [G2]. The equation of $\mathcal{E}_{\mathbb{Z}/6\mathbb{Z}}$ is given [GS2, Table 1]. We recall that the reducible fibers of the fibration $\mathcal{E}_{\mathbb{Z}/6\mathbb{Z}}$ are $2I_6 + 2I_3 + 2I_2 + 2I_1$ (we will number the reducible fibers assuming that the first two reducible fibers are of type I_6, the third and the fourth are of type I_3, the fifth and the sixth are of type I_2). Let t_1 be a 6-torsion section, then $t_1 \cdot C_1^{(j)} = 1$ for all $j = 1, \ldots, 6$ and $t_1 \cdot C_i^{(j)} = 0$ if $i \neq 1$. Here we give two orthogonal copies of $E_8(-1)$ which are contained in $NS(X_{\mathbb{Z}/6\mathbb{Z}})$:

$$\begin{array}{ccccccccccccc}
C_2^{(3)} & - & C_0^{(3)} & - & s & - & C_0^{(1)} & - & C_5^{(1)} & - & C_4^{(1)} & - & C_3^{(1)} \\
 & & & & | & & & & & & & & \\
 & & & & C_0^{(5)} & & & & & & & &
\end{array}$$

$$\begin{array}{ccccccccccccc}
C_2^{(4)} & - & C_1^{(4)} & - & t_1 & - & C_1^{(2)} & - & C_2^{(2)} & - & C_3^{(2)} & - & C_4^{(2)} \\
 & & & & | & & & & & & & & \\
 & & & & C_1^{(6)} & & & & & & & &
\end{array}$$

There exists a Morrison–Nikulin involution, ι_M, of the surface which switches them (cf. [Mo, Proposition 5.7]). This involution switches the singular fibers of the same type and acts on $MW(\mathcal{E}_{\mathbb{Z}/6\mathbb{Z}})$ sending $s \leftrightarrow t_1$, $t_2 \leftrightarrow t_5$, $t_3 \leftrightarrow t_5$ (this is deduced by the fact that $t_i \cdot C_j^{(h)} = \iota_M(t_i) \cdot \iota_M(C_j^{(h)})$). We observe that the action of ι_M on the sections is $t_i \leftrightarrow t_{-i+1}$. Hence ι_M is the involution defined in proof of Proposition 8.1. In particular $\mathcal{D}_6 = \langle \iota_M, \sigma_6 \rangle$.
The case $G = (\mathbb{Z}/3\mathbb{Z})^2$. The elliptic fibration $\mathcal{E}_{(\mathbb{Z}/3\mathbb{Z})^2}$ has 8 fibers of type I_3 (cf. [GS2, Table 2]). The transcendental lattice of the surface $X_{(\mathbb{Z}/3\mathbb{Z})^2}$ is isometric to $U(3) \oplus U(3)$. The automorphisms ι switches pairwise of fibers of type I_3. On the quotient elliptic fibration this gives 4 fibers of type I_3. Moreover the automorphism ι fixes two smooth fibers, and on each of these fibers it has four fixed points. On the quotient elliptic fibration $\mathcal{E}/\iota$ this gives two fibers of type I_0^*. So the trivial lattice of the elliptic fibration $\underbrace{\mathcal{E}/\iota\ \text{is}}$ $U \oplus A_2^{\oplus 4} \oplus D_4^{\oplus 2}$ and there are no sections of infinite order, because $rk(Tr_{\mathcal{E}/\iota}) = rk(NS(\widetilde{X_{(\mathbb{Z}/3\mathbb{Z})^2}/\iota}))$. The discriminant of the trivial lattice is $3^4 2^4$. By [Shim] the elliptic fibration $\mathcal{E}/\iota$ is such that $tors(MW(\mathcal{E}_{(\mathbb{Z}/3\mathbb{Z})^2/\iota})) = \{1\}$. This implies that

$d(NS(\widetilde{X_{(\mathbb{Z}/3\mathbb{Z})^2}/\iota})) = 3^4 2^4 = 6^4$. Moreover it is easy to find the discriminant form of $NS(\widetilde{X_{(\mathbb{Z}/3\mathbb{Z})^2}/\iota})$. The discriminant form of the transcendental lattice is the opposite of the one of the Néron–Severi group. Hence one can directly check that the hypothesis of Lemma 8.3 with $n = 6$ are satisfied by $T_{\widetilde{X_{(\mathbb{Z}/3\mathbb{Z})^2}/\iota}}$, and so there exist a lattice L such that $T_{\widetilde{X_{(\mathbb{Z}/3\mathbb{Z})^2}/\iota}} = L(6)$. The discriminant form of L, computed by the one of $T_{\widetilde{X_{(\mathbb{Z}/3\mathbb{Z})^2}/\iota}}$, is trivial, hence L is a unimodular lattices with signature $(2,2)$. This implies that $L \simeq U \oplus U$ and so $T_{\widetilde{X_{(\mathbb{Z}/3\mathbb{Z})^2}/\iota}} \simeq U(6) \oplus U(6)$. Since $U(6) \oplus U(6) \simeq (T_{X_{(\mathbb{Z}/3\mathbb{Z})^2}})(2)$, the involution ι is a Morrison–Nikulin involution (cf. [Mo, Theorems 5.7, 6.3]).

The case $G = \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$. The elliptic fibration $\mathcal{E}_{\mathbb{Z}/2\mathbb{Z}\times\mathbb{Z}/4\mathbb{Z}}$ has 4 fibers of type I_4 and 4 fibers of type I_2 (cf. [GS2, Table 2]). The automorphisms ι switches two fibers of type I_4, two other fibers of type I_4, two fibers of type I_2 and the remaining two fibers of type I_2. Moreover it fixes two smooth fibers and in particular four distinct points on each of these fibers. As in case $G = (\mathbb{Z}/3\mathbb{Z})^2$ this gives the quotient elliptic fibration. The trivial lattice of the elliptic fibration $\mathcal{E}/\iota$ is $U \oplus A_1^{\oplus 2} \oplus A_3^{\oplus 2} \oplus D_4^{\oplus 2}$ and there are no sections of infinite order. The discriminant of the trivial lattice is 2^{10}. By [Shim] the elliptic fibration $\mathcal{E}/\iota$ is such that $tors(MW(\mathcal{E}_{\mathbb{Z}/2\mathbb{Z}\times\mathbb{Z}/4\mathbb{Z}}/\iota)) = (\mathbb{Z}/2\mathbb{Z})^2$. This implies that $d(NS(\widetilde{X/\iota})) = 2^{10}/2^4 = 2^6$. We observe that also $d(NS(X_{\mathbb{Z}/2\times\mathbb{Z}/4\mathbb{Z}})) = 2^6$. In particular this implies that ι is not a Morrison–Nikulin involution, indeed, if $Y = \widetilde{X/i}$ with a Morrison–Nikulin involution i, then $T_Y = T_X(2)$ (cf. [Mo, Theorems 5.7, 6.3]) so in particular $d(NS(X)) = 2^{22-\rho(X)} d(NS(Y))$, where $\rho(X)$ is the Picard number of X. □

Proposition 8.5. *Let* G, ι *and* $\mathcal{E}_G$ *be as in Proposition 8.1. Let* $K = \langle G, \iota\rangle$. *Then:*

i) *the desingularization,* $\widetilde{X_G/K}$, *of the quotient* X_G/K *is such that* $NS(\widetilde{X_G/K}) \simeq NS(\widetilde{X_G/\iota})$;

ii) *the desingularization of the quotient* $\widetilde{X_G/K}$ *is such that* $NS(\widetilde{X_G/K})$ *is an overlattice of index 4 of* $U(2) \oplus M_K$.

If $G = \mathbb{Z}/n\mathbb{Z}$, $n = 5, 6$, *then* $T_{\widetilde{X_G/\iota}} \simeq T_{\widetilde{X_G/K}} \simeq U(2) \oplus U(2n)$, *if* $G = (\mathbb{Z}/3\mathbb{Z})^2$, *then* $T_{\widetilde{X_G/\iota}} \simeq T_{\widetilde{X_G/K}} \simeq U(6) \oplus U(6)$.

Proof. Here we consider the case $n = 6$. The others are very similar.

Proof of i). Let $\mathcal{E}/\iota$ be the quotient elliptic fibration. The involution ι switches the two fibers of type I_6, the two fibers of type I_3, the two fibers of type I_2. On $\mathcal{E}/\iota$ these reducible fibers correspond to one fiber of type I_6, one of type I_3, one of type I_2. On the fibration $\mathcal{E}$, ι fixes two smooth fibers. On these fibers ι fixes 4 points (the points P such that $2P = T$ where T is the intersection of the fiber with the section t_1). Hence these two fibers corresponds on $\mathcal{E}/\iota$ to 2 fibers of type I_0^*. So the trivial lattice of the fibration $\mathcal{E}/\iota$ is $U \oplus A_5 \oplus A_2 \oplus A_1 \oplus D_4^{\oplus 2}$. Choosing a numbering of the fibers and of the irreducible components $(D_i^{(j)})$ of the fibers we assume that the first fiber is of type I_6, the second of type I_3, the third of type I_2, the fourth and the fifth of type I_0^* and $F \simeq D_0^{(4)} + D_1^{(4)} + 2D_2^{(4)} + D_3^{(4)} + D_4^{(4)} \simeq D_0^{(5)} + D_1^{(5)} + 2D_2^{(5)} + D_3^{(5)} + D_4^{(5)}$ where F is the class of the fiber and $'\simeq'$ is the linear equivalence of divisors.

The elliptic fibrations with trivial lattice $U \oplus A_5 \oplus A_2 \oplus A_1 \oplus D_4^{\oplus 2}$ admit a 2-torsion section r (cf. [Shim]) and up to a choice of numbering of the components of the reducible fibers we can assume, by the height formula, that the intersection numbers are:

$$r \cdot D_3^{(1)} = r \cdot D_0^{(2)} = r \cdot D_1^{(3)} = r \cdot D_1^{(4)} = r \cdot D_1^{(5)} = 1$$

and the other intersections are zero.

Now let us consider the quotient $X/\mathcal{D}_6$. Under $\mathcal{D}_6$, the orbit of the curve $C_0^{(1)}$ consists of all the curves $C_i^{(j)}$, $i = 0, \ldots, 5$, $j = 1, 2$, so it is made up of all the components of the two fibers of type I_6. Hence the image of these two fibers on $X/\mathcal{D}_6$ is a nodal curve, where the node is the image of the singular points of the two fibers of type I_6. Analogously, the image of the two fibers of type I_3 is a nodal curve. The node is the image of the singular points of these fibers, and it is a singularity of type A_1 for the surface $X/\mathcal{D}_6$. Indeed on X the singular points of the fibers of type I_3 are fixed by $\sigma_{t_1}^2$ and hence their stabilizer is $\mathbb{Z}/2\mathbb{Z}$, this implies that the corresponding point on the quotient is a singularity of type A_1. With the same argument, one notices that the two fibers of type I_2 correspond on the quotient to a nodal curve, whose node is a singular point for $X/\mathcal{D}_6$ of type A_2 and the two fibers of type I_1 correspond on the quotient to a nodal curve, whose node is a singular point for $X/\mathcal{D}_6$ of type A_5. Each of the two smooth fibers fixed by ι correspond on the quotient to a rational curves with 4 singularities of type A_1. Blowing up these singularities we find the quotient elliptic fibration $\mathcal{E}/\mathcal{D}_6$. Its singular fibers are$I_1+I_2+I_3+I_6+2I_0^*$. The trivial lattice of this fibration is $U\oplus A_1\oplus A_2\oplus A_5\oplus D_4^{\oplus 2}$. It has a zero section (the image of the sections of $\mathcal{E}$), it has no sections of infinite order (because it has Picard number equal to 18, as the one of X and its trivial lattice has rank 18) and has a 2-torsion section ([Shim]).The singular fibers of the $\mathcal{E}/\mathcal{D}_6$ are the same as the ones of $\mathcal{E}/\iota$ and $MW(\mathcal{E}/\mathcal{D}_6)=MW(\mathcal{E}/\iota)$. Moreover up to a choice of the numbering on the components of the reducible fibers one can assume that the intersection property of the 2-torsion section on $\mathcal{E}/\mathcal{D}_6$ are the same as the one of the 2-torsion section of $\mathcal{E}/\iota$. The Néron–Severi group of an elliptic surface is completed determined by its singular fibers and by its sections and hence $NS(\widetilde{X/\iota})\simeq NS(\widetilde{X/\mathcal{D}_6})$.

Proof of ii). Let us consider again the quotient $\widetilde{X/\mathcal{D}_6}$. We will call the components of its reducible fibers $B_i^{(j)}$ and we number the fibers assuming that the first is of type I_6, the second of type I_2, the third of type I_3 and the fourth and fifth of type I_0^*. By the construction described above the components which do not intersect the zero section of the fibers of type I_6, I_3, I_2 arise from the desingularization of singular points of type A_5, A_2, A_1 respectively on the quotient $X/\mathcal{D}_6$. So the components $B_i^{(1)}$, $i=1,\ldots,5$, $B_j^{(2)}$, $j=2,3$ and $B_1^{(3)}$ are curves in $M_{\mathcal{D}_6}$. The fibers of type I_0^* on $\widetilde{X/\mathcal{D}_6}$ are obtained as desingularization of a rational curve which has 4 singularity of type A_1, hence the components $B_i^{(j)}$ of the fourth and fifth fibers which resolve the singularity of type A_1 on the quotient $X/\mathcal{D}_6$ are $B_h^{(k)}$, $h=0,1,3,4$, $k=4,5$ and hence they are curves in $M_{\mathcal{D}_6}$. The curves $B_i^{(1)}$, $i=1,\ldots,5$, $B_j^{(2)}$, $j=2,3$, $B_1^{(3)}$, $B_h^{(k)}$, $h=0,1,3,4$, $k=4,5$ generates the lattice $A_5\oplus A_2\oplus A_1^{\oplus 9}$ which is a sublattice of $M_{\mathcal{D}_6}$. The lattice $M_{\mathcal{D}_6}$ is described in [X, Table 2, Lemma 2] as overlattice of $A_5\oplus A_2\oplus A_1^{\oplus 9}$ obtained by adding two 2-divisible classes and hence $A_5\oplus A_2\oplus A_1^{\oplus 9}\hookrightarrow M_{\mathcal{D}_6}$ is an inclusion with index 4. Indeed the classes

$$\begin{array}{ll} v:=\frac{1}{2}(B_0^{(4)}+B_1^{(4)}+B_3^{(4)}+B_4^{(4)}+B_0^{(5)}+B_1^{(5)}+B_3^{(5)}+B_4^{(5)}) & \sim(F+B_2^{(4)}+B_2^{(5)}),\\ w:=\frac{1}{2}\left(B_1^{(1)}+B_3^{(1)}+B_5^{(1)}+B_1^{(3)}+\sum_{j=4}^{5}\left(B_3^{(j)}+B_4^{(j)}\right)\right) & \sim r-2F-s, \end{array}$$

where r is the 2-torsion section of $\mathcal{E}/\mathcal{D}_6$, are in $M_{\mathcal{D}_6}\subset NS(\widetilde{X/\mathcal{D}_6})$.

The orthogonal lattice to $M_{\mathcal{D}_6}$ in $NS(\widetilde{X/\mathcal{D}_6})$ is made up of the two classes F and $z:=-2s+B_1^{(4)}+2B_2^{(4)}+B_3^{(4)}+B_4^{(4)}+B_1^{(5)}+2B_2^{(5)}+B_3^{(5)}+B_4^{(5)}$. They generate a lattice which is isometric to $U(2)$. Hence $U(2)\oplus M_{\mathcal{D}_6}\hookrightarrow NS(\widetilde{X}/\mathcal{D}_6)$ with a finite index. This index is $4=(d(U(2)\oplus M_{\mathcal{D}_6}))\,/\left(d(NS(\widetilde{X}/\mathcal{D}_6)\right)$ and in fact the classes

$$\begin{array}{ll} z+B_1^{(4)}+B_3^{(4)}+B_4^{(4)}+B_1^{(5)}+B_3^{(5)}+B_4^{(5)} & \in U(2)\oplus M_{\mathcal{D}_6}\\ F+B_0^{(4)}+B_1^{(4)}+B_3^{(4)}+B_4^{(4)} & \in U(2)\oplus M_{\mathcal{D}_6} \end{array}$$

are divisible by 2 in $NS(\widetilde{X}/\mathcal{D}_6)$.

Let ν be a Morrison–Nikulin involution on a K3 surface S. Then $T_{\widetilde{S/\nu}}\simeq T_S(2)$ (cf. [Mo, Theorems 5.7 and 6.3])). The transcendental lattice of $X_{\mathbb{Z}/n\mathbb{Z}}$ is computed in [GS1] and is isometric to $U\oplus U(n)$. So $T_{\widetilde{X_{\mathbb{Z}/n\mathbb{Z}}/\iota}}\simeq U(2)\oplus U(2n)$. Since $NS(\widetilde{X/\iota})\simeq NS(\widetilde{X/K})$, the discriminant form of $T_{\widetilde{X/\iota}}$ is isometric to the one of $T_{\widetilde{X/K}}$. The lattice with such a discriminant form is uniquely determined, up to isometry, indeed by Lemma 8.3 it is multiple of a certain lattice which is uniquely determined by [N2, Corollary 1.13.3]. Hence $T_{\widetilde{X_{\mathbb{Z}/n\mathbb{Z}}/\iota}}\simeq T_{\widetilde{X_{\mathbb{Z}/n\mathbb{Z}}/\mathcal{D}_6}}$. □

Remark 8.6. As in Remark 7.9 one obtains that for $n=5,6$ the lattice $DIH_{2n}(16)$ described in [GL] is isometric to the lattice $\Omega_{\mathcal{D}_n}(-1)$, and, by Remark 6.3, they are both isometric to $MW(F_{gen}^{(n)})$, described in [Shio4]. The isometry $\Omega_{\mathcal{D}_5}(-1)\simeq DIH_{10}(16)$ was proved also in [G2]. □

In Proposition 4.1 we exhibit K3 surfaces obtained as quotient of other K3 surfaces X with an abelian group G of symplectic automorphisms, such that $NS(X) \simeq U \oplus M_G$. In the case $G = \mathcal{D}_5$, $\mathcal{D}_6$, $\mathfrak{A}_{3,3}$, $\mathbb{Z}/2\mathbb{Z} \times \mathcal{D}_4$ this is not possible, indeed there exists no K3 surfaces with Néron–Severi group isometric to $U \oplus M_G$ for one of these group G.

9. Final remarks on finite groups of symplectic automorphisms.

In Section 8 we analyzed the following property of a pair of finite groups (H, G):

(P)
- G acts symplectically on a K3 surface;
- H is a subgroup of G;
- the K3 surfaces admitting H as group of symplectic automorphisms admits necessary the group G as group of symplectic automorphisms.

In particular we analyzed this property by a geometrical point of view, constructing elliptic surfaces and using them to show that certain pairs (H, G) satisfy property (P). Here we reconsider this property from a point of view more related to the lattices.

Proposition 9.1. *A pair of groups (H, G) satisfy property (P) if and only if $H \subset G$ and $rk\Omega_H = rk\Omega_G$.*

Proof. If (P) holds, then $\Omega_H \subseteq \Omega_G$ (because $H \subset G$ and the K3 surfaces which admits G as group of symplectic automorphisms admits H as group of symplectic automorphisms), in particular $\mathrm{rk}\Omega_H \leq \mathrm{rk}\Omega_G$. If X is a generic algebraic K3 surface such that H acts symplectically on X, then $\rho(X) = 1 + rk\Omega_H$. If Property (P) holds, then $\Omega_K \subset NS(X)$ (c.f. Proposition 1.1). Since Ω_G is negative definite, $rk\Omega_G \leq \rho(X) - 1 = rk\Omega_H$, hence $\mathrm{rk}\Omega_H = \mathrm{rk}\Omega_G$.
Viceversa if $H \subset G$, then $\Omega_H \subseteq \Omega_G$. Since $\mathrm{rk}\Omega_H = \mathrm{rk}\Omega_G$, the inclusion $\Omega_H \hookrightarrow \Omega_G$ has a finite index. As both Ω_H and Ω_G are primitively embedded in the Néron–Severi group of the surface the index of the inclusion $\Omega_H \hookrightarrow \Omega_G$ is 1, which implies Property (P). □

We recall that for each group G of symplectic automorphisms on a K3 surface $\mathrm{rk}\Omega_G = \mathrm{rk}M_G$ and $\mathrm{rk}M_G$ is computed by [X].

Remark 9.2. There are no pairs (H, G) with the property (P) and $\mathrm{rk}\Omega_G < 16$. Indeed from Xiao's list we deduce that $\mathrm{rk}\Omega_G \in \{8, 12, 14, 15\}$. If G is such that $\mathrm{rk}\Omega_G = 8$, then $G = \mathbb{Z}/2\mathbb{Z}$. In the other case there is more then one group, but there are no inclusions among them. For example the groups G such that $\mathrm{rk}\Omega_G = 12$ are $\mathbb{Z}/3\mathbb{Z}$ and $(\mathbb{Z}/2\mathbb{Z})^2$, but of course there are no inclusions between them, and hence it is not possible to have a pair (H, G) with the property (P) and $\mathrm{rk}\Omega_G = 12$. The other cases are similar. □

Remark 9.3. Let us assume $\mathrm{rk}\Omega_G = 16$. The only pairs of groups (H, G) satisfying the property (P) are $(\mathbb{Z}/5\mathbb{Z}, \mathcal{D}_5)$, $(\mathbb{Z}/6\mathbb{Z}, \mathcal{D}_6)$, $(\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}, \mathbb{Z}/2\mathbb{Z} \times \mathcal{D}_4)$, $((\mathbb{Z}/3\mathbb{Z})^2, \mathfrak{A}_{3,3})$ (i.e. the pairs studied in Section 8). This follows by the list in [X]: the only groups G such that $\mathrm{rk}\Omega_G = 16$ are $\mathbb{Z}/5\mathbb{Z}$, $\mathcal{D}_5$, $\mathbb{Z}/6\mathbb{Z}$, $\mathcal{D}_6$, $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$, $\mathbb{Z}/2\mathbb{Z} \times \mathcal{D}_4$, $(\mathbb{Z}/3\mathbb{Z})^2$, $\mathfrak{A}_{3,3}$, $\mathfrak{A}_4$. So one has only to check that all the possible inclusions among these groups are those listed above. In particular the group $\mathfrak{A}_4$ is the only one which does not appear in a pairs (H, G). □

Remark 9.4. In Xiao's list there are 31 groups G acting symplectically on a K3 surface such that $\mathrm{rk}\Omega_G = 19$. By [Mu] there are 11 maximal groups acting symplectically on a K3 surface and for each of them the rank of the associated lattice Ω is 19. Hence there exist pairs (H, G) such that (H, G) satisfy the property (P) and $\mathrm{rk}\Omega_H = 19$. It would be interesting to find explicit examples to show this fact from a geometrical point of view. □

References

[BPV] W. Barth, C. Peters, A. van de Ven, *Compact complex surfaces*, Ergebnisse der Mathematik und ihrer Grenzgebiete 3. Springer-Verlag, Berlin, (1984).

[G1] A. Garbagnati, *Symplectic automorphimsms on Kummer surfaces,* eprint, arXiv:0802.0369v1 [math.AG].

[G2] A. Garbagnati, *The dihedral group $\mathcal{D}_5$ as group of symplectic automorphisms on K3 surfaces,* eprint, arXiv:0812.4518v1 [math.AG].

[GL] R. L. Griess Jr., C. H. Lam, *EE_8-lattices and dhiedral groups*, eprint, arXiv:0806.2753v1 [math.RT]

[GS1] A. Garbagnati, A. Sarti, *Symplectic automorphisms of prime order on K3 surfaces*, J. Algebra, **318** (2007), 323–350.

[GS2] A. Garbagnati, A. Sarti, *Elliptic fibrations and symplectic automorphisms on K3 surfaces*, to appear in Comm. in Alg., now avaible arXiv:08013992v2 [math.AG].

[vGS] B. van Geemen, A. Sarti, *Nikulin involutions on K3 surfaces*, Math. Z. **255**, no.4, (2007), 731–753.

[Kl] R. Kloosterman *Elliptic K3 surface with geometric Mordell–Weil rank 15*, Canad. Math. Bull. **50** (2007), 215–226.

[Ku] M. Kuwata *Elliptic K3 surfaces with a given Mordell–Weill rank*, Comment. Math. Univ. St. Paul. **49** (2000), 91–100.

[Mi] R. Miranda, *The basic theory of elliptic surfaces*, Università di Pisa, Dottorato di ricerca in matematica, ETS Editrice Pisa, (1989).

[Mo] D.R. Morrison, *On K3 surfaces with large Picard number*, Invent. Math. **75** (1986), 105–121.

[Mu] S. Mukai, *Finite groups of automorphisms of $K3$ surfaces and the Mathieu group*, Invent. Math. **94** (1988), 183–221.

[N1] V.V. Nikulin, *Finite groups of automorphisms of Kählerian $K3$ surfaces*, (Russian) Trudy Moskov. Mat. Obshch. **38** (1979), 75–137. English translation: Trans. Moscow Math.Soc. **38** (1980), 71–135.

[N2] V. V. Nikulin, *Integral symmetric bilinear forms and some applications*, Izv. Math. Nauk SSSR **43** (1979) 111–177, English transltion: Math. USSR Izvestija **14** (1980), 103–167.

[PS] I. Piatetski-Shapiro, I.R. Shafarevich, *Torelli's theorem for algebraic surfaces of type K3*, Izv. Akad. Nauk. SSSR Ser. Mat. **35** (1971), 530–572. English translation: Math. USSR, Izv. **5** (1971), 547–587.

[Shim] I. Shimada, *On elliptic K3 surfaces*, Michigan Math. J. 47 (2000), 423–446, (eprint version with the complete list arXiv:math/0505140v3 [math.AG]).

[Shio1] T. Shioda, *On Mordell–Weil lattice*, Comment. Math. Univ. St. Paul. **39** (1990), 211–240.

[Shio2] T. Shioda, *A note on K3 surfaces and sphere packings*, Proc. Japan Acad, **76A** (2000), 68–72.

[Shio3] T. Shioda, *The Mordell–Weil Lattice of* $y^2 = x^3 + t^5 - \frac{1}{t^5} - 11$, Comment. Math. Univ. St. Pauli, **56** (2007), 45–70.

[Shio4] T. Shioda, *K3 surfaces and sphere packing*, J. Math. Soc. Japan, **60** (2008), 1083–1105.

[ST] J. H. Silverman, J. Tate, *Rational points on elliptic curves*, Undergraduate Texts in Mathematics, Springer-Verlag, New York (1992).

[T] J.Top, *Descent by 3-isogeny and 3-rank of quadratic fields*, Advances in number theory (Kingston, ON, 1991), 303–317, Oxford Univ. Press, New York (1993).

[W] U. Whitcher, *Symplectic Automorphisms and the Picard Group of a K3 Hypersurface*, eprint, arXiv:09020601v1 [math.AG].

[X] G. Xiao, *Galois cover between K3 surfaces*, Ann. Inst. Fourier **46**, (1996), 73–88.

Alice Garbagnati, Dipartimento di Matematica, Università di Milano, via Saldini 50, I-20133 Milano, Italia
E-mail address: `alice.garbagnati@unimi.it`